

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
10 June 2004 (10.06.2004)

PCT


(10) International Publication Number  
**WO 2004/047615 A2**

BEST AVAILABLE COPY

- (51) International Patent Classification<sup>7</sup>: **A61B**
- (21) International Application Number:  
PCT/US2003/037393
- (22) International Filing Date:  
21 November 2003 (21.11.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/428,050 21 November 2002 (21.11.2002) US
- (71) Applicant (for all designated States except US): **WYETH**  
[US/US]; Five Giralda Farms, Madison, NJ 07940 (US).
- (72) Inventor; and  
(75) Inventor/Applicant (for US only): **GREENBERGER,**  
Lee, M. [US/US]; \*.
- (74) Agents: **FEHLNER, Paul, F. et al.**; Darby & Darby P.C.,  
Post Office Box 5257, New York, NY 10150-5257 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **HEMIASTERLIN AFFINITY PROBES AND THEIR USES**

**Tubulin source:**



**Coomassie**

**Fluorography**

(57) Abstract: Photoaffinity probes are provided that are based hemiasterlin and derivative compounds thereof. Use of these probes to identify binding sites for these and other drugs, particularly anti-tubulin drugs, are also provided as are methods for identifying new drugs (*e.g.*, new anti-tubulin drugs) that bind to these binding sites.

WO 2004/047615 A2

2200063-WO0

## HEMIASTERLIN AFFINITY PROBES AND THEIR USES

5

### 1. CROSS-REFERENCE TO RELATED APPLICATIONS

Priority is claimed under 35 U.S.C. 119(e) to copending U.S. provisional patent application Serial No. 60/428,050 filed on November 21, 2002. The contents of this prior application are hereby incorporated by reference and in their entirety.

10

### 2. FIELD OF THE INVENTION

The present invention relates to methods and compositions for identifying anticancer drugs and, in particular, for identifying binding sites and/or targets for anticancer drugs. In particular, the invention provides photoaffinity probes that mimic the binding of hemiasterlin derivatives, including the hemiasterlin derivative HTI-286, to tubulin. The invention also relates to methods for using such probes – including methods for identifying drug binding sites on tubulin, as well as diagnostic and prognostic methods that use these probes to identify cells containing mutant tubulin such as tumor cells. The invention additionally relates to methods using target binding sites that are identified with such probes; *e.g.*, to identify new binding compounds and potential therapeutic compounds, and/or to identify potentially drug resistant cells and tumors.

15

20

### 3. BACKGROUND OF THE INVENTION

$\alpha$ - and  $\beta$ -tubulin heterodimers polymerize to form microtubules which are vital for mitosis, motility, secretion, and proliferation (Rowinsky and Tolcher, in *Cancer Principles and Practice* (Devita *et al.*, eds.) 6<sup>th</sup> Ed. 2001, 431-452). Agents that bind tubulin and disrupt the function of microtubules are of great interest as some of these agents are routinely used to treat cancer. Three well-defined classes of drugs that bind tubulin have been previously identified: (1) colchicine, (2) vinca alkaloids and (3) taxanes (Hamel, *Med. Res. Rev.* 1996, 16:207-231). Colchicine and vinca alkaloids bind to distinct sites in tubulin, prevent the formation or extensions of microtubules, and therefore induce the depolymerization of microtubules (Hamel, *Med. Res. Rev.* 1996, 16:207-231; Downing, *Annu. Rev. Cell Dev. Biol.* 2000, 16:89-111). Photoaffinity-labeling and electron microscopy studies have revealed that taxanes bind to a distinct site

25

30

35

within  $\beta$ -tubulin (Downing, *Annu. Rev. Cell Dev. Biol.* 2000, 16:89-111). At stoichiometric amounts in relation to tubulin, taxanes stabilize microtubules and therefore actually enhance polymerization (Jordan, *Curr. Med. Chem. Anti-Canc. Agents* 2002, 2:1-17). However, at low concentrations, which are sufficient to inhibit cell division, all  
5 antimicrotubule agents alter microtubule dynamics without causing marked depolymerization or polymerization effects (Jordan, *Curr. Med. Chem. Anti-Canc. Agents* 2002, 2:1-17).


Vinca alkaloids and taxanes such as paclitaxel and docetaxel have been widely used to treat solid tumors. However, resistance to paclitaxel and vinca alkaloids is easily  
10 demonstrated in tissue culture systems and occurs frequently either at the onset or during the course of multiple cycles of chemotherapy in patients (Rowinsky and Tolcher, in *Cancer Principles and Practice* (Devita *et al.*, eds.) 6<sup>th</sup> Ed. 2001, 431-452). In tissue culture systems, vinca alkaloids, paclitaxel, and docetaxel, are excellent substrates for the ABC drug efflux pump, P-glycoprotein. This protein can be overexpressed in tumor cells  
15 in response to chemotherapeutic drugs and is believed to mediate resistance to these agents.

Recently a fourth class of anti-tubulin compounds, known as hemiasterlins, has been described. See, for example, Talpir *et al.*, *Tetrahedron Lett.* 1994, 35:4453-4456; Gambel *et al.*, *Bioorg. Med. Chem.* 1999, 7:1611-1615; Coleman *et al.*, *Tetrahedron*  
20 1995, 51:10653-10662; and Anderson *et al.*, *Cancer Chemother. Pharmacol.* 1997, 39:223-226. Hemiasterlins, which are tripeptide compounds isolated from marine sponges, induce microtubule depolymerization, cell cycle arrest and ultimately cell death (Anderson *et al.*, *Cancer Chemother. Pharmacol.* 1997, 39:223-226; Talpir *et al.*, *Tetrahedron Lett.* 1994, 35:4453-4456; Hamel and Covel, *Curr. Med. Chem. Anti-Canc. Agents* 2002, 2:19-53). The use of hemiasterlin compounds in cancer therapy has also  
25 been described. See, for example, International Patent Publication Nos. WO 99/32509 and WO 96/33211. See also, U.S. Patent No. 6,153,590. Methods for obtaining hemiasterlin compounds have additionally been described, both by isolating the compounds from marine sponges (U.S. Patent Nos. 5,661,175 and 6,153,590) and by  
30 chemical synthesis (Anderson & Coleman, *Tetrahedron Lett.* 1997, 38:317-320).

Synthetic hemiasterlin analogs and derivative compounds have also been described (see, for example, International Patent Publication No. WO 99/32509) and these compounds also have cytotoxic and anti-mitotic activity. In particular, provisional

U.S. patent application Serial Nos. 60/411,883 and 60/493,841 filed on September 20, 2002 and August 8, 2003, respectively, describe various hemiasterlin derivative compounds, including one compound known as HTI-286.

HTI-286, which has the chemical structure set forth in Formula I, below, has a weak interaction with P-glycoprotein and has also been shown to overcome resistance of other anti-tubulin drugs, such as taxanes, both *in vitro* and in xenograft tumor models (Loganzo *et al.*, *Cancer Res.* 2003, 63:1838-1845). Clinical trials of HTI-286 in cancer patients are in progress (Ratain *et al.*, *Proc. Am. Soc. Clin. Onc.* 2003, abstract 516).



(Formula I)

While hemiasterlins and its analogs/derivatives represent a promising class of new anti-tubulin drugs, it is expected that resistance to this drug and other anti-microtubule agents will continue to be a problem. This is because resistance to antimicrotubule agents is multifactorial and is due to transporter pumps, alterations in tubulin, as well as changes in mechanisms that mediate cell death after microtubules have been disrupted (Dumontet *et al.*, *J. Clin. Oncol.* 1999, 17:1061-1070). Therefore, the identification of new antimicrotubule agents that overcome resistance mechanisms may be useful clinically.

In order to identify such agents an understanding of the intracellular drug interactions (for example, with tubulin) is necessary. Unlike other anti-tubulin compounds, however, the tubulin binding site for hemiasterlins and their analogs (*e.g.*, HTI-286) as well as any peptide-based inhibitors remains unknown and the exact mechanism(s) by which these compounds disrupt tubulin polymerization is poorly understood, at best (Hamel and Covell, *Curr. Med. Chem. Anti-Canc. Agents* 2002, 2:19-53). Knowledge of the binding site and mechanism(s) of action would greatly facilitate the design and identification of new, more effective, anti-tubulin compounds, as well as mutations (such as in  $\alpha$ - and/or  $\beta$ -tubulin) that are associated with resistance to hemiasterlins. Hence, there continues to exist a need for anti-tubulin compounds, including new hemiasterlin derivatives or other peptide-like inhibitors, that are effective for inhibiting cancer and tumor cell growth. There also exists a need for methods (including diagnostic and prognostic methods) for identifying tumor cells and, in

particular, for identification cell lines and/or mutations that are resistant to such hemiasterlin compounds.


\* \* \* \* \*

5           The citation and/or discussion of a reference in this section and throughout the specification is provided merely to clarify the description of the present invention and is not an admission that any such reference is "prior art" to the invention described herein.

#### 10           4. SUMMARY OF THE INVENTION

10           The present invention at least partly overcomes the above-described problems by providing compounds that are useful, *e.g.*, for detecting hemiasterlin binding sites. In addition, the invention also provides binding data from such probes that can be used, *inter alia*, to design novel hemiasterlin compounds. Such hemiasterlin compounds are themselves useful, *e.g.*, as new anti-cancer drugs. Binding data obtained using  
15       compounds of the invention can also be used to identify cells (particularly cancer cells) that are resistant to or are likely to be resistant to hemiasterlin drugs. Such methods therefore are also provided in the present invention.


          The invention therefore provides, more specifically, compounds represented by the chemical formula:



20

          or a pharmaceutically acceptable salt thereof, in which at least one of the substituents A, B, E, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and/or R<sub>9</sub> comprises a photoreactive group such as a benzophenone or an azide moiety and in which the substituents A, B, E, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and R<sub>9</sub>, when not a photoreactive moiety, is as described in the Sections below.

25           Preferred compounds of the invention have the chemical formula:



or a pharmaceutically acceptable salt thereof in which  $R_1$  is either a photoreactive moiety or an aryl moiety,  $R_2$  is either a photoreactive moiety, and alkyl moiety of H, and wherein at least one of  $R_1$  and  $R_2$  comprises a photoreactive moiety.

Particularly preferred compounds of the invention are:

5


4-benzoyl-N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-N1-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N1,3-dimethyl-L-valinamide;

N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-

10

N, $\beta$ , $\beta$ -trimethyl-L-phenylalaninamide;

and pharmaceutically acceptable salts thereof. Preferred compounds of the invention also include ones having the chemical formula:



15

and pharmaceutically acceptable salts thereof.

The invention also provides methods that use such compounds to identify tubulin binding sites, *e.g.*, for hemiasterlin drugs. Such methods generally provide contacting a compound of the invention (*e.g.*, one of the compounds described, *supra*) to a sample containing tubulin such that the compound can irreversibly bind tubulin in the sample. The tubulin is then separated into a plurality of different tubulin fragments, and at least one tubulin fragment is identified that has the compound bound thereto. Those fragments

that have a compound of the invention bound to them are thereby identified as fragments that comprise a tubulin binding site.

The invention additionally provides methods for identifying hemiasterlin competitors – *i.e.*, compounds that compete with a hemiasterlin compound for binding to  
5 tubulin. Such compounds can themselves be useful, *e.g.*, as novel anti-tubulin and anti-cancer drugs. The methods generally involve contacting both a test compound and a probe, which is one of the compounds of the invention (*e.g.*, any of the compounds described *supra*), is contacted to a sample that contains tubulin, and binding of the probe  
10 to tubulin is detected. This binding is compared to binding of the probe to tubulin that is observed in the absence of the test compound. Where binding in the presence of the test compound is lower than in the absence of the test compound, then the test compound is identified as a hemiasterlin inhibitor.

In various aspects of these different embodiments, the compound of the tubulin sample is preferably irradiated (*e.g.*, with UV-light) after being contacted with the probe  
15 compound, *e.g.*, to promote irreversible, covalent binding of the probe compound to tubulin. The probe compound is also preferably labeled so that its binding to tubulin (or to a tubulin fragment) can be detected by detecting the label. In other aspects of these methods, tubulin in the sample can be separated, *e.g.*, by chemical digest (for example, with formic acid or CNBr) or by enzymatic digestion (for example, with Lys C, Trypsin  
20 or subtilisin).

These and other particular embodiments of the invention are described in detail in the following Sections.

## 5. BRIEF DESCRIPTION OF THE DRAWINGS

25 **Figure 1A** illustrates results of purified bovine brain tubulin (2.5  $\mu\text{M}$ ) and purified Hela cell tubulin (10  $\mu\text{M}$ ) incubated for 30 minutes at room temperature with 2.5  $\mu\text{M}$  (3.6  $\mu\text{Ci}$ ) of [ $^3\text{H}$ ]-probe 1. After incubation, the samples were irradiated for 2 hours at 4°C with 360 nm UV light. Proteins were then resolved by SDS-PAGE and analyzed by Coomassie blue staining (left) or fluorography (right) of the gel as described below.  
30 **Figure 1B** illustrates the labeling of KB-3-1 cell lysates (50  $\mu\text{g}$ ) (lanes 1 and 2) and purified bovine brain (BB) tubulin (5  $\mu\text{g}$ ) (lanes 3 and 4) incubated for 30 minutes at room temperature with 2.5  $\mu\text{M}$  [ $^3\text{H}$ ]-probe 1. Samples in lanes 1 and 3 were pre-incubated with 250  $\mu\text{M}$  non-radiolabeled probe 1 prior to incubation with radiolabeled

materials. Samples were analyzed as described in **Figure 1A**. Lanes marked "M" are molecular weight markers in kDa.

**Figure 2** illustrates the inhibition of  $\alpha$ -tubulin labeling by Probe 1 in the presence of HTI-286. The gel image resulted from the following procedure: Bovine brain tubulin (2.5  $\mu$ M) was incubated without or with 1 mM HTI-286 for 15 min at 4°C, followed by addition of increasing concentrations of [ $^3$ H]-probe 1 (0.025-2.5  $\mu$ M) (0.036-3.6  $\mu$ Ci). After incubation for 30 minutes at 37°C, samples were irradiated with 360 nm UV light for 30 minutes and separated by SDS-PAGE. Gels were exposed to film by fluorography.

**Figures 3A-3C** provide data for the digestion of Probe 1-photolabeled tubulin by formic acid. Photolabeled tubulin was resolved by SDS-PAGE under conditions that allowed  $\alpha$ - and  $\beta$ -tubulin to comigrate. The tubulin band was excised from the gel and digested with 75% formic acid at 37°C. After 44 or 72 hours, formic acid digestion products were separated on a 10-20% Tris-tricine gel. **Figure 3A** shows a diagram of the predicted formic acid cleavage sites and cleavage products for  $\alpha$ -tubulin. **Figures 3B and 3C** are visualizations of the gel fragments stained by Coomassie Blue and of the radioactive fragments detected by fluorography, respectively.

**Figures 4A-4D** provide data for the digestion of Probe 1-photolabeled tubulin with trypsin. **Figure 4A** is a diagram demonstrating known cleavage sites within  $\alpha$ -tubulin after native tubulin digestion by trypsin. **Figures 4B and 4C** are images of trypsin cleavage fragments observed after Coomassie Blue staining or fluorography methods, respectively. For these images, tubulin (15  $\mu$ g) was incubated with 0.25  $\mu$ M (0.36  $\mu$ Ci) of [ $^3$ H]-Probe 1 for 30 minutes at room temperature and irradiated at 4°C with 360 nm UV light. After 2 hours, the samples were digested with 0.8  $\mu$ g trypsin at 30°C for 0 to 30 minutes. After stopping the reaction with 0.01 mM leupeptin, samples were resolved on 10-20% Tris-tricine gels. Alternatively, as demonstrated in **Figure 4D**, 1 mm gel slices were removed from the gel described in **Figure 4C** and quantitated to provide a radioactivity profile of the length of the gel.

**Figures 5A-5D** provide data for the digestion of Probe 1 photolabeled tubulin by CNBr. **Figures 5A and 5B** show images of the CNBr-digested tubulin fragments visualized by either fluorography or silver staining, respectively. These gel images were



produced as follows: After native tubulin was photolabeled as described in Section 6.1.5,  $\alpha$ - and  $\beta$ -tubulin were visualized in gels by Coomassie blue staining. Protein within each species was extracted, reduced, alkylated, and digested with 150 mg/mL CNBr in 70% formic acid at 37°C. After 48 hours, samples were resolved by SDS-PAGE. **Figure 5C** diagrams the predicted position of the 7 kDa labeled peptide fragment from  $\alpha$ -tubulin obtained after CNBr digestion. **Figure 5D** shows all predicted CNBr fragments of rat  $\alpha$ -tubulin. The boxes indicate sequences that are confirmed using mass spectrometry.

**Figures 6A-6C** provide data for the digestion of tubulin by subtilisin before or after photoaffinity labeling with Probe 1. **Figure 6A** diagrams the predicted subtilisin cleavage site and cleavage products for  $\alpha$ -tubulin. **Figure 6B** provides Coomassie staining (top) and fluorographic (bottom) images of fragments photolabeled with Probe 1 and exposed to subtilisin as follows: Tubulin (5  $\mu$ g) was incubed without (Lanes 1 and 4) or with 0.2  $\mu$ g subtilisin for 60 minutes at 30°C and then photolabeled with 0.25  $\mu$ M [ $^3$ H]-Probe 1 (Lane 2) according to conditions specified in Section 6.1.11. Alternatively, photolabeling was done before subtilisin digestions (Lane 3). In both cases samples were then separated by SDS-PAGE. **Figure 6C** shows Coomassie staining and fluorographic images of tubulin fragments prepared as follows: tubulin (5  $\mu$ g) was pre-incubated with 0, 100, or 250  $\mu$ M non-radiolabeled probe 1 for 15 minutes, labeled with [ $^3$ H]-Probe 1 as described above, and then digested with 0.2  $\mu$ g subtilisin for 60 minutes at 30°C as described in Section 6.1.11. Again, samples were separated by SDS-PAGE.

**Figures 7A-7B** present data for the photolabeling of bovine brain tubulin (Lane 1) and HeLa cell tubulin (Lane 2) using Probe 2 as described in Sections 6.1.5 and 6.6.1. **Figure 7A** provides an image of a 7.5% Tris-HCl gel of bovine brain and HeLa cell tubulin stained with Coomassie blue. **Figure 7B** shows an autoradiograph of the gel in **Figure 7A**.

**Figures 8A-8E** present data for trypsin, subtilisin, formic acid, and Lys C digestions of purified bovine brain tubulin before or after photolabeling with Probe 2. Methods for these procedures are described in Sections 6.1.7, 6.1.12, and 6.1.13. **Figure 8A** and **8B** provide Coomassie blue stained gel and fluorographic images, respectively, of tubulin fragments from trypsin, subtilisin, and formic acid digestions. **Figure 8B** shows a fluorographic image of fragments that have been radiolabeled with

Probe 2. Similarly, **Figure 8C** and **8D** provide Commassie blue stained gel and fluorographic images, respectively, of tubulin fragments from Lys C digestions. **Figure 8D** shows a fluorographic image of fragments that have been radiolabeled with Probe 2. **Figure 8E** diagrams the predicted Lys C fragments (described by their molecular weight, length, and sequence) of rat  $\alpha$ -tubulin. The 12.8 kDa fragment is in bold.

**Figures 9A-9B** provide data for the digestion of Probe 2-labeled bovine brain tubulin using CNBr. Probe 2 labeled tubulin was separated into  $\alpha$  and  $\beta$ -tubulin subunits by SDS-PAGE and separately digested with CNBr. **Figure 9A** displays a silver stained 4-20% Tris-Tricine gel of tubulin fragments after CNBr digestion. **Figure 9A** shows an audioradiograph containing radiolabeled fragments prepared according to Section 6.1.8.

**Figure 10** displays Commassie blue stained 7.5% Tris-HCl SDS-PAGE gels of purified bovine brain tubulin incubated with 50  $\mu$ M drug and digested with subtilisin for various times. Reactions were quenched and fragments were separated according to Section 6.1.14.

**Figure 11** is a structural model depicting the  $\alpha\beta$ -tubulin dimer. The bracketed areas show the approximate regions for the  $\alpha$ - and  $\beta$ -tubulin subunits. The general area of the Probe 1 and Probe 2 binding regions are identified with arrows. The bound GDP, GTP and paclitaxel are shown as ball and stick models. (PDB accession number 1JFF).

**Figure 12** shows a protein sequence alignment for amino acid residues 1-451 of human (*Homo sapien*, designated htub1 and htub2), primate (*Macaca fascicularis* and *Macaca mulatta*, designated ptub1 and ptub2 respectively), mouse (*Mus musculus*, designated motub1), hamster (*Cricetulus griseus*, designated hamtub1), rat (*Rattus norvegicus*, designated rattub1), chicken (*Gallus gallus*, designated chicktub1), and frog (*Xenopus laevis*, designated frogtub1)  $\alpha$ -tubulin (Stachi *et al.*, *Biochem. Biophys. Res. Commun.* 2000, 270:1111-1118). The EMBL Accession Numbers for htub1, htub2, ptub1, ptub2, motub1, hamtub1, rattub1, chicktub1, and frogtub1 are AJ245922, K00558, X04757, AF141923, AJ245923, M12329, V01227, M16030, and X07046, respectively.

## 6. DETAILED DESCRIPTION

### 6.1. AFFINITY PROBES


The present invention provides compounds that are generally referred to here as  
5 “affinity probes” and which are useful to detect and identify binding sites for drugs –  
particularly anti-cancer drugs. In particular, affinity probes of this invention are useful to  
detect and identify binding sites for a class of anti-cancer drugs referred to here as “anti-  
tubulin” drugs or, alternatively, as “tubulin inhibitors” or “tubulin binding compounds.”  
Without being limited to any particular theory or mechanism of action, tubulin inhibitors  
10 are believed to exert their therapeutic effects by specifically binding tubulin (which can  
be either  $\alpha$ -tubulin or  $\beta$ -tubulin) and disrupting the polymerization of these subunits into  
microtubules in cells. Hence, affinity probes of this invention are useful for identifying  
binding sites of tubulin inhibitor drugs on tubulin.

Preferred affinity probes of the invention are analogs of a particular class of anti-  
15 tubulin drugs generally referred to as hemiasterlin derivatives or “hemiasterlins.”  
Hemiasterlins are natural products derived from marine sponges that induce microtubule  
depolymerization, cell cycle arrest and ultimately cell death (Anderson *et al.*, *Cancer  
Chemother. Pharmacol.* 1997, 39:223-226; Telpin *et al.*, *Tetrahedron Letters* 1994,  
35:4453-4456). The use of hemiasterlin compounds in cancer therapy has also been  
20 described. See, for example, International Patent Publication Nos. WO 99/32509 and  
WO 96/33211. See also, U.S. Patent No. 6,153,590. Methods for obtaining hemiasterlin  
compounds have additionally been described, both by isolating the compound from  
marine sponges (U.S. Patent Nos. 5,661,175 and 6,153,590) and by chemical synthesis  
(Anderson & Coleman, *Tetrahedron Letters* 1997, 38:317-320).

25 Synthetic hemiasterlin analogs and derivative compounds have also been  
described (see, for example, International Patent Publication No. WO 99/32509) and  
these compounds also have cytotoxic and anti-mitotic activity. Accordingly, such  
hemiasterlin derivative and analogue compounds are also considered hemiasterlins, at  
least for the purposes of describing and claiming this invention. It is to be understood  
30 that, for the purposes of describing this invention, the terms “hemiasterlin derivative” and  
“hemiasterlin analog” are used interchangeably. Both these terms (in all their variants)  
therefore generally refer to compounds that are derived from and/or are chemical analogs  
of a natural hemiasterlin compound.

Particularly preferred classes of hemiasterlin derivatives have been described in provisional U.S. patent application Serial Nos. 60/411,883 and 60/493,841 filed on September 20, 2002 and August 8, 2003, respectively. These include a particular hemiasterlin derivative known as HTI-286, which has the chemical structure set forth in

5 Formula I, below (see also Table I in the examples, *infra*).



Formula I

HTI-286 has a weak interaction with P-glycoprotein and has been shown to overcome resistance of other anti-tubulin drugs (*e.g.*, taxane) both *in vitro* and in xenograft tumor models (Loganzo *et al.*, *Cancer Res.* 2003, 63:1838-1845). The compound is therefore a particularly useful as an anti-cancer drug, and clinical trial of

10 HTI-286 in cancer patients are in progress (Ratain *et al.*, *Proc. Am. Assoc. Cancer Res.* 2003, abstract 516). Hence, affinity probes of the present invention that are analogs of HTI-286 are particularly preferred.


Accordingly, the present invention provides compounds that are hemiasterlin derivatives as set forth in U.S. provisional patent application Serial Nos. 60/411,883 and

15 60/493,841 filed on September 20, 2002 and August 8, 2003 respectively in which one or more chemical moiety or substituent is either replaced by or additionally comprises a photoreactive group. Examples of preferred photoreactive groups which may be present in a compound of the invention include an azide moiety and a benzophenone moiety, with

20 benzophenone moieties being preferred. Other photoreactive groups include, but are not limited to, diazo groups, diazirines, enones, sulfur radicals, halogenated substrates, nitrobenzenes, dizonium salts, sulfonium salts, as well as those groups cited by Fleming (*Tetrahedron* 1995, 51:12479-12520).

For example, the present invention provides compounds represented by

25 Formula II, below:



(Formula II)

wherein at least one of the substituents A, B, E, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and/or R<sub>9</sub> comprises a


photoreactive group such as an azide or a benzophenone moiety. In preferred embodiments, at least one of the substituents A, B, E, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and/or R<sub>9</sub> is a benzophenone moiety.

In Formula II, above, A can comprise a photoreactive group and/or is selected from the group consisting of an alkyl moiety of 1 to 10 carbon atoms, an alkenyl moiety of 2 to 10 carbon atoms, an aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms are optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, -Br, -Cl, -F, CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, NO<sub>2</sub>, -SO<sub>3</sub>H, SOR<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub> wherein R<sub>10</sub> is an alkyl moiety of 1 to 10 carbon atoms, an alkenyl moiety of 2 to 10 carbon atoms, and aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, aryl-R- or heteroaryl-R; or A can be OR, S(O)R, S(O)<sub>2</sub>R, SO<sub>2</sub>NR<sub>2</sub>, NR<sub>1</sub>R<sub>2</sub> or N<sub>3</sub>.

R can comprise a photoreactive group and/or is selected from the group consisting of H, an alkyl moiety of 1 to 18 carbon atoms, an alkenyl moiety of 2 to 18 carbon atoms, an aryl and a cyclic hydrocarbon moiety of 3 to 18 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms are optionally substituted with =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, -Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub> or -SO<sub>2</sub>R<sub>10</sub>; wherein R<sub>10</sub> is an alkyl moiety of 1 to 10 carbon atoms, an alkenyl moiety of 2 to 10 carbon atoms, and aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, aryl-R- or heteroaryl-R.

B is O or H<sub>2</sub>.

E can comprise a photoreactive group and/or is the moiety:



Alternatively, E can comprise a photoreactive group and/or comprises an aryl moiety, a 5 to 14-membered monocyclic, bicyclic or tricyclic saturated or unsaturated hydrocarbon ring moiety wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms wherein the carbon atoms

may optionally be substituted with: R, =O, =S, -OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SO<sub>2</sub>CR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -I, -Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub> or -SO<sub>2</sub>R<sub>10</sub>;

- 5 R<sub>1</sub> can comprise a photoreactive group and/or a moiety is selected from the group consisting of H, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S,  
 10 OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SO<sub>2</sub>CR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

- R<sub>2</sub> can comprise a photoreactive group and/or a moiety selected from the group consisting of H, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon  
 15 atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SO<sub>2</sub>CR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -  
 20 CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

R<sub>1</sub> and R<sub>2</sub> together may optionally form a ring of 3 to 7 carbon atoms wherein carbon atoms may optionally be replaced with 0 to 2 nitrogen atoms, 0 to 2 oxygen atoms and 0 to 2 sulfur atoms;

- R<sub>3</sub> can comprise a photoreactive group and/or a moiety selected from the group  
 25 consisting of H, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SO<sub>2</sub>CR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -  
 30 NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

R<sub>4</sub> can comprise a photoreactive group and/or a moiety selected from the group consisting of H, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon

atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

R<sub>3</sub> and R<sub>4</sub> together may optionally form a ring of 3 to 7 carbon atoms wherein carbon atoms may optionally be replaced with 0 to 2 nitrogen atoms, 0 to 2 oxygen atoms and 0 to 2 sulfur atoms;

R<sub>5</sub> can comprise a photoreactive group and/or a moiety selected from the group consisting of H, OH, NHR, SH, aryl, heteroaryl, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

R<sub>5</sub> and A may optionally form a ring of 5 to 7 carbon atoms wherein carbon atoms may optionally be replaced with 0 to 2 nitrogen atoms, 0 to 2 oxygen atoms, and 0 to 2 sulfur atoms;


R<sub>6</sub> can comprise a photoreactive group and/or a moiety selected from the group consisting of H, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

R<sub>7</sub> can comprise a photoreactive group and/or a moiety selected from the group consisting of H, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0


to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

5 R<sub>8</sub> can comprise a photoreactive group and/or a moiety selected from the group consisting of H, an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms may optionally be substituted with: =O, =S,  
10 OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>;

R<sub>9</sub> can comprise a photoreactive group and/or a moiety selected from the group consisting of:



wherein W' can comprise a photoreactive group and/or is selected from the group consisting of SO<sub>2</sub>R<sub>16</sub>, SO<sub>3</sub>R<sub>14</sub>, SO<sub>2</sub>NR<sub>14</sub>R<sub>15</sub>, P(O)(OR<sub>14</sub>)(OR<sub>15</sub>), CN, OH, tetrazole, a moiety



20 and SO<sub>2</sub>NRR where the R groups may form a 4 to 8 membered ring wherein the carbon atoms may optionally be replaced with 0 to 2 nitrogen atoms, 0 to 2 oxygen atoms and 0 to 2 sulfur atoms.


R<sub>14</sub> and R<sub>15</sub> may comprise a photoreactive moiety and/or are independently selected from the group consisting of H and an alkyl moiety of 1 to 6 carbon atoms. R<sub>16</sub>  
25 is an alkyl moiety of 1 to 6 carbon atoms in length. D is O or OH and/or may comprise a photoreactive group.

Z and Y may optionally form a ring of 5 to 7 carbon atoms wherein the carbon atoms may optionally be replaced by 0 to 2 nitrogen atoms, 0 to 2 oxygen atoms and/or 0




to x sulfur atoms. Y may comprise a photoreactive moiety and/or may comprise an alkyl moiety of 1 to 10 carbon atoms optionally substituted with R, ArylR-, or X or an alkenyl moiety of 2 to 10 carbon atoms optionally substituted with R, ArylR- or X. Z may comprise a photoreactive moiety and/or may comprise a moiety selected from the group consisting of: H, an alkyl moiety of 1 to 6 carbon atoms, -NRN(R)<sub>2</sub>, R, aryl, heteroaryl, aralkyl, -OR, -SH, -SR, -NH<sub>2</sub>, -NHR, -NROR, -N(R)<sub>2</sub>, NH-NH<sub>2</sub>, and NRR; where the R group may form a 4 to 8 membered ring wherein carbon atoms may optionally be replaced with 0 to 2 nitrogen atoms, 0 to 2 oxygen atoms and/or 0 to 2 sulfur atoms, -NHCH(R<sub>11</sub>)COOH; and -NRCH(R<sub>11</sub>)COOH in which R<sub>11</sub> may comprise a photoreactive group and/or is a moiety having the formula R or -(CH<sub>2</sub>)<sub>n</sub>NR<sub>12</sub>R<sub>13</sub> wherein n=1-4 and both R<sub>12</sub> and R<sub>13</sub> are independently selected from the group consisting of H, R and -C(NH)(NH<sub>2</sub>).

Alternatively, Z may comprise a photoreactive moiety and/or may comprise a moiety having selected from:




wherein the dotted line indicates an optional chemical bond, Q is (CH<sub>2</sub>)<sub>m</sub>, G<sub>1</sub> is selected from the group consisting of O, N and S, m is an integer of between 1 and 3, and u is an integer of between 0 and 5. R<sub>17</sub> is a phenyl or O-(CH<sub>2</sub>)<sub>n</sub>phenyl, R<sub>18</sub> is H or OH, and R<sub>19</sub> is selected from a chemical bond, an alkyl moiety of 1 to 10 carbon atoms optionally substituted with an alkyl moiety of 1 to 10 carbon atoms and alkoxy of 1 to 10 carbon atoms.

R<sub>20</sub> is selected from OR<sub>14</sub>, NH-R<sub>21</sub>, a moiety of the formula:



and a moiety of the formula:




$R_{21}$   $R_{22}$  are each preferably an alkyl moiety of 1 to 10 carbon atoms optionally substituted with aryl and/or heteroaryl moieties.

5 X is defined as a moiety selected from the group consisting of: -OH, -OR, =O, =S, -O<sub>2</sub>CR, -SH, -SR, -SOCR, -NH<sub>2</sub>, -NHR, -N(R)<sub>2</sub>, -NHCOR, -NRCOR, -I, -Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R, -CHO, -COR, -CONH<sub>2</sub>, -CONHR, -CON(R)<sub>2</sub>, -COSH, -COSR, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR and -SO<sub>2</sub>R.

10 In preferred embodiments the invention provides a compound according to Formula II, as defined above, wherein at least one of the substituents  $R_3$ ,  $R_4$ ,  $R_5$ , and  $R_7$  comprises a photoreactive group such as an azide moiety or a benzophenone moiety, with a benzophenone moiety being particularly preferred. In particularly preferred embodiments a compound according to Formula II comprises a photoreactive group (preferably a benzophenone moiety) at the substituent  $R_5$  or at the substituent  $R_7$ .

15

The present invention also provides compounds represented by Formula III, below:



(Formula III)

20 wherein at least one of the substituents  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$  and/or  $R_9$  comprises a photoreactive group such as an azide or a benzophenone moiety. In preferred embodiments, at least one of the substituents  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$  and/or  $R_9$  is a benzophenone moiety.

25 In a compound of Formula III, above,  $R_1$  can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>,

-I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; and aryl-R-.

R<sub>2</sub> can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; and aryl-R-.


Alternatively, R<sub>1</sub> and R<sub>2</sub> taken together with the nitrogen atom to which they are attached can be three to seven membered ring.

R<sub>3</sub> can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; and aryl-R-.

R<sub>4</sub> can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; and aryl-R-.

Alternatively, R<sub>3</sub> and R<sub>4</sub> taken together with the carbon to which they are attached can form a three to seven membered ring.

R<sub>5</sub> can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, , CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; aryl-R- and aryl. In a preferred embodiment R<sub>5</sub> is an idoly moiety of the formula:



in which R<sub>17</sub> is H, a photoreactive group or an optionally substituted acyl group. R<sub>18</sub>, Q<sub>1</sub>, Q<sub>2</sub>, Q<sub>3</sub> and Q<sub>4</sub> are independently selected from H, a photoreactive group, a halogen, alkyl, acyl, -OH, -O-alkyl, -O-acyl, -NH<sub>2</sub>, -NH-alkyl, -N(alkyl)<sub>2</sub>, -NH-acyl, -NO<sub>2</sub>, -SH, -S-alkyl and -S-acyl in which the alkyl and acyl groups of the substituents are optionally substituted.


R<sub>6</sub> can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; and aryl-R-.

R<sub>7</sub> can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear,

branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; and aryl-R-.

R<sub>8</sub> can comprise a photoreactive group and or may comprise a moiety selected from the group consisting of: H; a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms, and zero to four sulfur atoms, said carbon atoms being optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CH-O, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, or -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group; and aryl-R-.

R<sub>9</sub> can comprise a photoreactive group and/or the moiety:



In the above formula, R is preferably a saturated or unsaturated moiety having a linear, branched, or cyclic skeleton containing one to ten carbon atoms, zero to four nitrogen atoms, zero to four oxygen atoms and zero to four sulfur atoms. The carbon atoms in R can be optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R<sub>10</sub>, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub> wherein R<sub>10</sub> is a linear, branched or cyclic, one to ten carbon saturated or unsaturated alkyl group.

X may be a photoreactive group and/or comprises a moiety selected from the group consisting of: -OH, -OR, =O, =S, -O<sub>2</sub>CR, -SH, -SR, -SOCR, -NH<sub>2</sub>, -NHR, -N(R)<sub>2</sub>, -NHCOR, -NRCOR, -I, -Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>R, -CHO, -COR, -CONH<sub>2</sub>, -CONHR, -CON(R)<sub>2</sub>, -COSH, -COSR, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR, and -SO<sub>2</sub>R.


Y may be a photoreactive group and/or comprises a moiety selected from the group consisting of: a linear, saturated or unsaturated on to six carbon alkyl group that is optionally substituted with R, ArylR-, or X.

Z may be a photoreactive group and/or comprises a moiety selected from the group consisting of -OH, -OR, -SH, -SR, -NH<sub>2</sub>, -NHR, -N(R)<sub>2</sub>, -NHCH(R<sub>11</sub>)COOH. R<sub>11</sub> may comprise a photoreactive group and/or is a moiety having the formula R or - (CH<sub>2</sub>)<sub>n</sub>NR<sub>12</sub>R<sub>13</sub>, where n is an integer of between 1 and 4, and where R<sub>12</sub> and R<sub>13</sub> are independently selected from the group consisting of H, R, -C(NH) (NH<sub>2</sub>) and pharmaceutically acceptable salts thereof.

In preferred embodiments the invention provides a compound according to Formula II, as defined above, wherein at least one of the substituents R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, and R<sub>7</sub> comprises a photoreactive group such as an azide moiety or a benzophenone moiety, with the benzophenone moiety being particularly preferred. In particularly preferred embodiments a compound according to Formula II comprises a photoreactive group (preferably a benzophenone moiety) at the substituent R<sub>5</sub> or at the substituent R<sub>7</sub>.

In particularly preferred embodiments the invention provides compounds that are derivatives of the hemiasterlin analog HTI-286 of Formula I, *supra*, that additionally comprise a photoreactive group and/or in which a photoreactive group is optionally substituted for one or more of the substituent moieties in Formula I. Exemplary photoreactive groups include an azide and a benzophenone moiety, with the benzophenone moiety being particularly preferred.

For example, compounds of the invention that have the chemical structure set forth in Formula IV, below, are particularly preferred:



(Formula IV)

25

In Formula IV, above, the substituents R<sub>1</sub> and R<sub>2</sub> are independently selected provided that at least one of the substituents R<sub>1</sub> and R<sub>2</sub> comprises a photoreactive moiety such as a benzophenone or azide moiety. R<sub>1</sub> can be a photoreactive moiety or an aryle moiety such as a benzene ring. R<sub>2</sub> can be a photoreactive moiety, and alkyl moiety or H.


Specific preferred compounds of the invention include the compounds:

4-benzoyl-N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-N1-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N1,3-dimethyl-L-valinamide;

N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N, $\beta$ , $\beta$ -trimethyl-L-phenylalaninamide;


and pharmaceutically acceptable salts thereof.

Specific preferred compounds of the invention include compounds, referred to here as Probe 1 and Probe 2, respectively, with the formula:



(Probe 1),

and



(Probe 2).

It is understood that, when describing compounds of the present invention, the term alkyl means a saturated linear or branched hydrocarbon moiety of between 1 and 20 carbon atoms, with hydrocarbon moieties of 1 to 10 carbon atoms being preferred.. In some embodiments of the invention the alkyl moiety may optionally be 1 to 18 carbon atoms or 1 to 6 carbon atoms. Examples include, but are not limited to, Examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-

methylbutyl, 2,2-dimethylpropyl, n-hexyl, 2-methylphenyl, 2,2-dimethylbutyl, n-heptyl, 2-methylhexyl, and the like unless otherwise specified. The carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms, and 0 to 4 sulfur atoms, and the carbon atoms are optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SO<sub>2</sub>CR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, aryl-R- and heteroaryl-R.

10 The term alkenyl refers to an unsaturated linear or branched hydrocarbon moiety, preferably between about 2 to 10 carbon atoms and containing at least one carbon-carbon double bond. Each double bond in an alkenyl moiety is independently a cis, trans or nongeometric isomer wherein carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms and 0 to 4 sulfur atoms. In addition, the carbon atoms are optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SO<sub>2</sub>CR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, aryl-R- and heteroaryl-R.

The term cyclic hydrocarbon moiety refers to a saturated or unsaturated cyclic hydrocarbon moiety (preferably of 3 to 10 carbon atoms) as well as to a monocyclic cycloalkyl or cyclalkenyl ring of 3 to 10 carbon atoms. Carbon atoms in a cyclic hydrocarbon moiety may optionally be replaced with 0 to 4 nitrogen atoms, 0 to 4 oxygen atoms and 0 to 4 sulfur atoms. In addition, the carbon atoms are optionally substituted with: =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SO<sub>2</sub>CR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is a an alkyl moiety of 1 to 10 carbon atoms, alkenyl moiety of 2 to 10 carbon atoms, aryl and a cyclic hydrocarbon moiety of 3 to 10 carbon atoms, aryl-R- and heteroaryl-R.

In some embodiments of the invention the cyclic hydrocarbon may optionally be a 5 to 14-membered monocyclic, bicyclic or tricyclic saturated or unsaturated hydrocarbon ring moiety wherein the carbon atoms may optionally be replaced with 0 to 4 nitrogen atoms,



0 to 4 oxygen atoms and 0 to 4 sulfur atoms, wherein the carbon atoms may optionally be substituted with R, =O, =S, OH, -OR<sub>10</sub>, -O<sub>2</sub>CR<sub>10</sub>, -SH, -SR<sub>10</sub>, -SOCR<sub>10</sub>, -NH<sub>2</sub>, -NR<sub>10</sub>H, -N(R<sub>10</sub>)<sub>2</sub>, -NHCOR<sub>10</sub>, -NR<sub>10</sub>COR<sub>10</sub>, -I, Br, -Cl, -F, -CN, -CO<sub>2</sub>H, -CHO, -COR<sub>10</sub>, -CONH<sub>2</sub>, -CONHR<sub>10</sub>, -CON(R<sub>10</sub>)<sub>2</sub>, -COSH, -COSR<sub>10</sub>, -NO<sub>2</sub>, -SO<sub>3</sub>H, -SOR<sub>10</sub>,  
5 or -SO<sub>2</sub>R<sub>10</sub> where R<sub>10</sub> is as defined above.

The term aryl refers to an aromatic hydrocarbon moiety having 6, 10 or 14 carbon atoms and preferably having between 6 and 10 carbon atoms optionally substituted with R, X or Z (as defined *supra*).

The term heteroaryl refers to a 5- or 6-membered heterocyclic ring, which can be  
10 fused to another 5- or 6-membered heterocyclic ring, especially heteraromatic rings that contain 1 to 3 heteroatoms selected which can be, *e.g.*, O, N or S optionally substituted with R or X or fused to a cyclic hydrocarbon moiety of 3 to 10 carbon atoms. Examples of preferred heteroaromatic rings include but are not limited to: thienyl, furyl, indolyl, pyrrolyl, thiophenyl, benzofuryl, benzothiophenyl, quinolyl, isoquinolyl, imidazolyl,  
15 thiazolyl, oxazolyl and pyridyl.

The term alkoxy refers to an alkyl-O- group in which the alkyl group is as defined above. Exemplary alkoxy groups include but are not limited to methoxy, ethoxy, n-propoxy, l-propoxy, n-butoxy and t-butoxy.

Aralkyl refers to an aryl-alkyl group in which the aryl and alkyl group are as  
20 defined above. Non-limiting examples of aralkyl groups include benzyl and phenethyl.

The term phenyl refers to a 6-membered carbon aromatic ring.

Preferably, the recitation of a compound of the invention (*e.g.*, a compound of Formula II or III as defined, *supra*) includes all possible salts of the compound, and also denotes all possible isomers possible within the structural formula specified for such  
25 compound, including geometrical and optical isomers. Unless otherwise stated, materials described herein comprising a compound for which isomers exist are to be regarded as covering individual isomers as well as mixtures of isomers, including racemic mixtures.

It is, moreover, to be understood that compounds of the invention may comprise on or more detectable labels, including one or more radiosotopes such as <sup>3</sup>H (*i.e.*,  
30 "tritium), <sup>13</sup>C, <sup>32</sup>P or <sup>35</sup>S to name a few. For instance, the Examples *infra* describe embodiments in which the benzephenone moiety in compounds of the invention comprises <sup>3</sup>H rather than normal <sup>1</sup>H.

## 6.2. USES OF AFFINITY PROBES

### 6.2.1 Identification of Tubulin Binding Sites

Affinity probes of the present invention are particularly well suited for identifying  
5 the binding site or sites for anti-tubulin compounds and, in particular, for identifying the  
binding site or sites of hemiasterlin compounds (including hemiasterlin derivatives and  
analogues). For instance, the Examples, *infra*, describe experiments that use affinity probes  
of the invention to identify hemiasterlin binding sites on tubulin. These experiments  
involve routine experimental techniques such as enzymatic digestion (for example,  
10 digestion using Lys C, Trypsin and/or subtilisin) as well as chemical digestion (*e.g.*, by  
formic acid and/or CNBr) to digest tubulin after irreversible binding to an affinity probe  
of the invention. For instance, in the Examples *infra*, a photoreactive group on the  
affinity probe is activated (*e.g.*, by UV-irradiation) after incubation with tubulin such that  
the reactive group irreversibly binds to the tubulin. Digestion fragments of tubulin can be  
15 subsequently analyzed to determine which fragment or fragments have an affinity probe  
bound thereto. The binding site or sites of the affinity probe (and hence of hemiasterlins)  
is then identified as the site or sites corresponding to the fragment or fragments with an  
affinity probe bound thereto.

As an example and not by way of limitation, digestion fragments can be separated  
20 (*e.g.*, by electrophoretic gels or other chromatography techniques known in the art) and a  
detectable label (for example, a radiolabel such as  $^3\text{H}$ ) on the affinity probe(s) can be  
detected. Alternatively, digestion fragments can be analyzed by mass spectroscopy to  
determine which fragment or fragments have an affinity probe bound thereto.

The Examples, *infra*, also identify particular binding sites on tubulin for anti-  
25 tubulin compounds and, in particular, for hemiasterlin compounds (including their  
derivatives and analogues). In particular, the results presented in these Examples  
demonstrate that hemiasterlins bind  $\alpha$ -tubulin in a region or regions between about amino  
acid residues 200 and 350 of  $\alpha$ -tubulin and, more preferably, between about amino acid  
residues 200-340.

30 In particular, the results in the Examples describe one embodiment (using the  
affinity probe referred to here as Probe 1) in which hemiasterlins bind to amino acid  
residues corresponding to a domain referred to as loop 8, helix 10 in  $\alpha$ -tubulin. In  
particular, this domain corresponds to a region comprising amino acid residues 300-350  
of  $\alpha$ -tubulin. More preferably, this region comprises amino acid residues 310-340, and

still more preferably comprises amino acid residues 314-338 of  $\alpha$ -tubulin. The Examples also describe another embodiment (using the affinity probe referred to here as Probe 2) in which hemiasterlins bind to a region comprising about amino acid residues 200-300 of  $\alpha$ -tubulin, and more preferably comprising amino acid residues 204-280 of  $\alpha$ -tubulin.

5 Those skilled in the art appreciate that these portions of the  $\alpha$ -tubulin amino acid sequences correspond to domains commonly referred to as helices 6-8, loop 7 and the N-terminal portion of the M-loop of  $\alpha$ -tubulin. It is understood, therefore, that hemiasterlins (including hemiasterlin derivatives and analogs such as HTI-286) can be expected to bind to either one or both of these regions on  $\alpha$ -tubulin.

10 It is well known that the sequence of  $\alpha$ -tubulin is highly conserved among different species of organisms and, in particular, among different species of vertebrates. See, in particular, the sequence alignment at **Figure 12**. Hence, while the experiments described in the Examples, *infra*, are done using bovine tubulin, it is understood that the results (including the binding domains and regions described above) are equally  
15 applicable to tubulin from other species of organisms, including tubulin from other vertebrate species such as those identified in **Figures 12**. Accordingly, the amino acid residues recited above and throughout this specification, unless otherwise noted, specifically refer to amino acid residues in the sequence of human  $\alpha$ -tubulin set forth in **Figure 12**. However, those skilled in the art will be able to readily identify corresponding  
20 domains and regions of other  $\alpha$ -tubulin amino acid sequences (including  $\alpha$ -tubulin sequences from other species of organisms, orthologs and homologs). For instance, residues of other  $\alpha$ -tubulin sequences shown in **Figure 12** that align with the amino acid sequences recited above are also understood to represent binding domains for hemiasterlin compounds.

25 Those skilled in the art can also readily align different  $\alpha$ -tubulin or other amino acid sequences, using routine algorithms such as FASTA (Pearson & Lipman, *Proc. Natl. Acad. U.S.A.* 1988, 85:2444-2448; Pearson, *Methods Enzymol.* 1990, 183:63-98), BLAST (Altschul *et al.*, *Nucl. Acids Res.* 1997, 25:3389-3402; Altschul, *J. Mol. Evol.* 1993, 36:290-300; Altschul *et al.*, *J. Mol. Biol.* 1990, 215:403-410), CLUSTAL and  
30 CLUSTALW (Higgins *et al.*, *Nucl. Acids. Res.* 1994, 22:4673-4680), to name a few. Generally, such alignment algorithms will be used with the standard or default parameters, including standard alignment scoring systems and/or a scoring matrix such as BLOSUM62. See, Henikoff & Henikoff, *Proc. Natl. Acad. Sci.* 1992, 89:10915-10919.

However, in certain circumstances that will be appreciated by those skilled in the art, it may be preferable to use nonstandard parameters and/or scoring matrices. For example, in embodiments where very similar amino acid sequences are being compared (such as sequences of  $\alpha$ -tubulin) it may be preferable to use a scoring matrix such as BLOSUM90,  
5 that has higher cutoffs.

#### 6.2.2 Assays for Characterizing Anti-Tubulin Compounds

Affinity probes of this invention can also be used to characterize compounds based on their ability to inhibit affinity probe binding and/or covalent labeling to tubulin.  
10 For example, if a compound significantly inhibits binding of an affinity probe, this compound can be characterized as being similar in structure, mechanism of action, and/or the site of binding compared to the probe. Alternatively, if a compound does not inhibit the binding of an affinity probe, this compound can be characterized as being different in structure, mechanism of action, and/or the site of binding compared to the probe.  
15 Compounds characterized according to such methods can possess inhibiting activity ranging anywhere from completely inhibiting an affinity probe from binding tubulin (*i.e.*, complete inhibition) to completely allowing the affinity probe to bind (*i.e.*, no inhibition). Compounds characterized according to these methods can also be partial inhibitors that, while not completely inhibiting binding of the affinity probe to tubulin, nevertheless do  
20 inhibit binding to some measurable extent.

Classes of compounds that may be used to inhibit probe binding include, but are not limited to, small molecules (*e.g.*, organic or inorganic molecules which are less than about 2 kDa in molecular weight, are more preferably less than about 1 kDa in molecular weight, and/or are able to cross the blood-brain barrier and affect tubulin or activities  
25 associated therewith) as well as macromolecules (*e.g.*, molecules greater than about 2 kDa in molecular weight). Compounds used to inhibit binding may also include peptides and polypeptides. Examples of such compounds (including peptides) include but are not limited to: soluble peptides; fusion peptide members of combinatorial libraries (such as ones described by Lam *et al.*, *Nature* 1991, 354:82-84; and by Houghten *et al.*, *Nature*  
30 1991, 354:84-86); members of libraries derived by combinatorial chemistry, such as molecular libraries of D- and/or L-configuration amino acids; phosphopeptides, such as members of random or partially degenerate, directed phosphopeptide libraries (see, *e.g.*, Songyang *et al.*, *Cell* 1993, 72:767-778); antibodies, including but not limited to polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies;

antibody fragments, including but not limited to Fab, F(ab')<sub>2</sub>, Fab expression library fragments, and epitope-binding fragments thereof.

General methods for determining in tubulin the inhibition of probe labeling by compounds are described here, and specific, non-limiting examples of these methods are demonstrated in the Examples, *infra* (see Sections 7.4.2 and 7.4.3). In general, a tubulin-containing sample (preferably comprising at least  $\alpha$ -tubulin) is incubated in the presence of both a test compound and a photoaffinity probe of the invention. In preferred embodiments, the tubulin sample is incubated first with the test compound, and the affinity probe is added to that sample after a time sufficient for the test compound to bind tubulin. However, the test compound and affinity probe can also be contacted to the tubulin sample concurrently, and incubated together for a time period sufficient for both compounds to bind. In preferred embodiments, the affinity probe is then covalently bound to the tubulin (for example, by irradiation the sample with UV light) to facilitate detection of the bound affinity probe. The tubulin sample can then be analyzed by routine methods, to determine whether and to what extent the affinity probe is bound to tubulin in the sample. Preferably, these results are compared to data from a control experiment, in which an affinity probe is incubated with the tubulin sample in the absence of a test compound.

Examples of assays that can be used to detect the affinity probe include, but are not limited to, detection of the affinity probe by radioactivity (*e.g.*, where a radiolabeled affinity probe is used) or fluorescence (*e.g.*, where a fluorescently labeled affinity probe is used). Alternatively, the affinity probe may also be detected by other methods such as mass spectroscopy or in an immunoassay (*e.g.*, with an antibody that specifically binds to the affinity probe or, alternatively, specifically binds to tubulin when an affinity probe is bound thereto).

The radioactivity of a probe can be detected by fluorography or scintillation counting as described in the Examples, *infra*. Probe fluorescence can be detected using methods well known in the art, such as fluorescence microscopy or fluorimetry.

In the use of antibodies, detecting the desired antibody can be accomplished by techniques known in the art, *e.g.*, radioimmunoassay, ELISA (enzyme-linked immunosorbant assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions, immunodiffusion assays, precipitation reactions, agglutination assays (*e.g.*, gel agglutination assays, hemagglutination assays),

complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc.

In one embodiment, antibody binding is assayed by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by assaying  
5 the binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

As demonstrated in Section 7.4.3, one preferred embodiment of this invention uses a radiolabeled photoaffinity hemiasterlin analog probe. In this example, known  
10 tubulin binding drugs, dolastatin-10, vinblastine, paclitaxel, and colchicine, are used as competitive inhibitors of probe binding to tubulin. However, it is understood that other compounds, which may or may not be known tubulin binding drugs, can also be characterized using such methods. In this Example, each test compound is preferably incubated with tubulin followed by the addition of probe. The probe can then be  
15 covalently attached to tubulin using UV light. Probe binding to tubulin is assayed, *e.g.*, by separating the samples on an SDS-PAGE gel, slicing the gel, and determining the radioactivity of each gel slice using scintillation counting. The amount of probe radioactivity in tubulin for each inhibition experiment is compared to probe radioactivity in tubulin in the absence of competitor. Using this method, dolastatin-10 and vinblastine  
20 inhibited probe labeling to a greater degree and, hence, are more closely related to the probe than paclitaxel or colchicine.

Results from such assays, which evaluate the ability of a test compound to inhibit hemiasterlin probes from labeling tubulin can also be compared to similar assays that use affinity probes for other classes of tubulin binding drugs. Such probes can mimic the  
25 binding of, for example, different classes of tubulin-binding drugs such as colchicine, vinca alkaloids, or taxanes. Based on the results of these inhibition studies compounds can be categorized. A compound, for example, can be considered most related in structure, mechanism of action, and/or site of binding to a particular probe if the degree of binding inhibition is higher than for other probes. Preferably, however, test compounds  
30 will show probe-labeling inhibition to at least one, all, none, or some (*e.g.* to varying degrees) of the probes.

### 6.2.3 Diagnostic and Prognostic Assays

Affinity probes of this invention are also useful for diagnostic and/or prognostic assays, *e.g.*, to identify cancerous cells and tissues both *in vitro* (such as in cell or tissue cultures) and *in vivo*. In particular, it is understood that tubulin binding compounds, including affinity probes of this invention, preferably target and bind to tubulin in cancerous cells and tissues which are rapidly dividing. Hence, such cells and tissues can be readily detected, by detecting binding of the affinity probes to tubulin in cells and tissue. Examples of tumors that can be detected according to the invention include sarcomas and carcinomas such as, but not limited to: fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

In other embodiments, affinity probes of the invention can also be used to identify cells and tumors to which a hemiasterlin compound (including hemiasterlin derivatives and analogs such as HTI-286) will bind, and which are therefore susceptible to treatment with those compounds. In other words, affinity probes of the invention can also be used to determine whether a particular cancer cell or tumor (for example, a cell or tumor obtained from a patient biopsy) are or will be resistant to treatment with hemiasterlin compounds such as HTI-286. Such assays are particularly useful, *e.g.*, for identifying therapeutic regimens that are effective for treating a particular cancer and/or for a particular patient.

Cells and/or tissue samples that are tested in such applications can be readily obtained – for example, by obtaining a biopsy sample from a patient who has or is suspected of having a cancer or tumor. Cells and tissue samples can also be maintained in culture, using routine culturing techniques, for later testing according to these methods.

The detection of tumors (including hemiasterlin susceptible tumors) in such assays is based on the the high binding affinity of the affinity probes to tubulin in such cell or tissue samples. Hence, high binding affinity of an affinity probe to cells or tissues is indicative that the cells and/or tissues are cancerous and, moreover, that they are

5 susceptible to treatment with a hemiasterlin compound. In preferred embodiments, cells or tissue samples can be exposed to probes by, for example, bathing in media containing the probe. Non-specific binding of the probe to the sample should be avoided by, for example, washing the sample with media without probe several times after specific binding has occurred. Probe specifically bound to tubulin can be detected using a variety

10 of methods including and of the methods described above – *e.g.*, by assaying the radioactivity and/or fluorescence of the probe, mass spectroscopy to determine the location of the probe, and/or antibody detection of the probe. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain Fab fragments, and Fab expression library. In addition, each of these antibodies may be fluorescently labeled.

15 The radioactivity of a probe can be detected by fluorography or scintillation counting as described in the Examples, *infra*. Probe fluorescence can be detected using methods well known in the art, such as fluorescence microscopy or fluorimetry.

In the use of antibodies, detecting the desired antibody can be accomplished by techniques known in the art, *e.g.*, radioimmunoassay, ELISA (enzyme-linked

20 immunosorbant assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions, immunodiffusion assays, *in situ* immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), precipitation reactions, agglutination assays (*e.g.*, gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, fluorescence

25 activated cell sorting (*i.e.* FACS), and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is assayed by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by assaying the binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in

30 an immunoassay and are within the scope of the present invention.

In another embodiment, probes described here may be used to detect (*e.g.*, to diagnose) dysproliferative changes (such as metaplasias and dysplasias) in epithelial tissues such as those in the cervix, esophagus, and lung. Thus, the present invention



provides for detection of conditions known or suspected of preceding progression to neoplasia or cancer, in particular, where non-neoplastic cell growth consisting of hyperplasia, metaplasia, or most particularly, dysplasia has occurred (for review of such abnormal growth conditions, see Robbins and Angell, 1976, *Basic Pathology*, 2d Ed.,  
5 W.B. Saunders Co., Philadelphia, pp. 68-79).

Hyperplasia is a form of controlled cell proliferation involving an increase in cell number in a tissue or organ, without significant alteration in structure or function. As but one example, endometrial hyperplasia often precedes endometrial cancer. Metaplasia is a form of controlled cell growth in which one type of adult or fully differentiated cell  
10 substitutes for another type of adult cell. Metaplasia can occur in epithelial or connective tissue cells. Atypical metaplasia involves a somewhat disorderly metaplastic epithelium. Dysplasia is frequently a forerunner of cancer, and is found mainly in the epithelia. It is the most disorderly form of non-neoplastic cell growth, involving a loss in individual cell uniformity and in the architectural orientation of cells. Dysplastic cells often have  
15 abnormally large, deeply stained nuclei, and exhibit pleomorphism. Dysplasia characteristically occurs where there exists chronic irritation or inflammation, and is often found in the cervix, respiratory passages, oral cavity, and gall bladder. For a review of such disorders, see Fishman et al., 1985, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia.

20 The methods described herein are not limited to diagnostic applications, but may also be used in prognostic applications, *e.g.*, to monitor the progression of a disease (such as cancer) that is associated with probe binding to tubulin, or to monitor a therapy thereto. Accordingly, prognostic methods of the invention may comprise, in one exemplary embodiment, monitoring probe binding of cells or tissues from an individual during the  
25 course of a treatment or therapy (for example, a drug treatment or chemotherapy regime) for cancer or for another disease associated with abnormal tubulin. Similarly, the methods of the invention may also be used to detect and identify diseased cells and tissue (for example, cancerous cells and tissue) during the course of a therapy.

### 30 6.3. HEMIASTERLIN BINDING DATA AND ITS USES

Information about binding sites and/or mechanisms of action for hemiasterlins and other anti-tubulin drugs is useful *inter alia* for identifying new tubulin binding compounds, including the identification of new hemiasterline derivatives and analogs, *e.g.*, by molecular modeling and/or structure activity relationship (SAR) studies. Hence,

the present invention provides methods for identifying new anti-tubulin drugs (particularly, new peptide-containing agents, such as hemiasterlins) which use SAR and/or molecular modeling to identify compounds that bind to one or more binding sites identified by an affinity probe of the invention. Information about binding sites can also  
5 be used to identify amino acid sequences and mutations (including tubulin amino acid sequences and mutations) that affect tubulin binding. Such mutations can give rise, *e.g.*, to tumor cells and/or cell lines that are resistant to certain hemiasterlins and other anti-tubulin compounds. Thus, the invention also provides methods for identifying such amino acid sequences, as well as diagnostic and prognostic methods for identifying cells  
10 that are resistant to (or are likely to be resistant to) certain hemiasterlins and/or other anti-tubulin compounds. These and other methods of the invention are described in detail herebelow.

#### 6.3.1 Identifying new tubulin binding compounds

15 Information about binding sites and/or mechanisms of action for hemiasterlins can be used, *e.g.*, in molecular modeling and/or structure activity relationship (SAR) studies to identify new anti-tubulin drugs and, in particular, to identify new hemiasterlin compounds (including new hemiasterlin derivatives and analogs). Such methods, which use information about binding sites obtained from affinity probes (see, the Examples  
20 *infra*) are therefore provided in the present invention.

For example, the present invention provides methods that use computer modeling algorithms and other techniques known in the art to identify compounds that bind (or are expected to bind) a hemiasterlin binding site identified by an affinity probe of this invention – such as one of the binding sites described above. Such methods generally use  
25 a three-dimensional or tertiary structure for tubulin or another drug target to model the binding site or sites of candidate compounds. The three dimensional structure of  $\alpha$ -tubulin, for example, has been determined and can be obtained, *e.g.*, from the Protein Data Bank (Bennan *et al.*, *Nucl. Acids Res.* 2000, 28:235-242) under the Accession Nos. 1TUB, 1JFF, and 1FFX. See also, Nogales *et al.*, *Nature* 1998, 391:199; Lowe *et al.*, *J.*  
30 *Mol. Biol.* 2001, 313:1045; and Gigart *et al.*, *Cell* 2000, 102:809.

Because  $\alpha$ -tubulin is highly conserved across different species of organisms (see, in particular, the alignment at **Figure 12**) the three dimensional structure of tubulin from one species of organism can be readily used to obtain a corresponding three-dimensional structure for tubulin from a different species of organism. In particular, it is understood

that the three-dimensional structure of all  $\alpha$ -tubulin molecules will be substantially similar regardless of the source or species of organism from which it is derived. In general, two three-dimensional structures are said to be substantially structurally similar to each other if their atomic coordinates have a root-mean square deviation (RMSD) less than or equal  
5 to about 1 angstrom, as calculated, *e.g.*, using the Molecular Similarity Module within the QUANT program (QUANTA, available from Molecular Simulations Inc., Sand Diego, CA).

Using routine computer modeling algorithms and other techniques that are well known in the art, interactions (*e.g.*, hydrogen bonding, hydrophobic, and/or electrostatic  
10 interactions) between hemiasterlin analogs of this invention and tubulin can be identified. Furthermore using these modeling techniques, a user may identify other compounds that are expected to bind to tubulin using similar interactions as the compounds described in this invention. More specifically, using the crystal structure of tubulin, those skilled in the art can identify compounds that bind by forming stabilizing interactions with tubulin,  
15 similar to the stabilizing interactions for hemiasterlin. Compounds identified using these modeling techniques can be expected to compete with hemiasterlin analogs for binding to tubulin.

Techniques and algorithms for such computer assisted molecular modeling are well known in the art. Exemplary programs that can be used in such methods include, but  
20 are not limited to: AutoDock (Morris *et al.*, *J. Computational Chem.* 1998, 19:1639-1662), FTDock (Katchelski-Katzin, *Proc. Natl. Acad. Sci.* 1992, 89:2195-2199; Gabb *et al.*, *J. Mol. Biol.* 1997, 272:106-120), GROMOS™ (van Gunstener & Berendsen, *Angew. Chem. Int. Ed. Engl.* 1990, 29:992-1023) and ICM (available from MolSoft LLC, San Diego CA) to name a few.

25 Classes of compounds that may be identified by such modeling techniques include, but are not limited to, small molecules (*e.g.*, organic or inorganic molecules which are less than about 2 kDa in molecular weight, are more preferably less than about 1 kDa in molecular weight, and/or are able to cross the blood-brain barrier and affect tubulin or activities associated therewith) as well as macromolecules (*e.g.*, molecules  
30 greater than about 2 kDa in molecular weight). Compounds identified by these modeling techniques may also include peptides and polypeptides. Examples of such compounds (including peptides) include but are not limited to: soluble peptides; fusion peptide members of combinatorial libraries (such as ones described by Lam *et al.*, *Nature* 1991,

354:82-84; and by Houghten *et al.*, *Nature* 1991, 354:84-86); members of libraries derived by combinatorial chemistry, such as molecular libraries of D- and/or L-configuration amino acids; phosphopeptides, such as members of random or partially degenerate, directed phosphopeptide libraries (see, *e.g.*, Songyang *et al.*, *Cell* 1993, 72:767-778); antibodies, including but not limited to polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies; antibody fragments, including but not limited to Fab, F(ab')<sub>2</sub>, Fab expression library fragments, and epitope-binding fragments thereof.

In other exemplary embodiments, compounds identified by computer modeling algorithms or other techniques that are well known in the art of this invention may actually have (or may be expected to have) improved binding or stabilizing interactions with tubulin. For example, compounds identified by these methods may form (or be expected to form) stronger and/or more specific hydrogen bonding interactions with tubulin. These compounds can be expected to be more potent in tubulin binding assays and functional assays that test a compounds ability to inhibit tubulin polymerization and cell proliferation.

*In vitro* or cell culture assays may also be used to determine whether a test compound functions as tubulin binder and an anti-microtubule agent. For instance, the Examples, *infra*, describe binding assays that determine the stability of binding between tubulin and test agents. Additionally, the Examples, *infra*, provide an assay to determine the ability of a compound to prevent tubulin polymerization *in vitro* and assay to assess cell proliferation in the presence of a test compound.

### 6.3.2 Identification of resistant cells and tumors

Information about binding sites and/or mechanisms of action for hemiasterlin compounds can also be used to identify cells and tissues (*e.g.*, tumors) that are either resistant or likely to be resistant to certain anti-tubulin drugs. In particular, such information can be used to identify cells and tissues that may be resistant to treatment with a hemiasterlin compound – including cells and tissues that are either resistant or are likely to be resistant to treatment with a hemiasterlin derivative or analog such as HTI-286.

As an example, and not by way of limitation, the Examples, *infra*, demonstrate that hemiasterlins bind  $\alpha$ -tubulin in a region or regions between about amino acid residues

200 and 350 of  $\alpha$ -tubulin, and more preferably between about amino acid residues 200-340.

In particular, the results in the below Examples describe one embodiment (using the affinity probe referred to here as Probe 1) in which hemiasterlins bind to amino acid  
5 residues corresponding to a domain referred to as loop 8, helix 10 in  $\alpha$ -tubulin. This domain corresponds to a region comprising amino acid residues 300-350 of  $\alpha$ -tubulin. More preferably, this region comprises amino acid residues 310-340, and still more preferably comprises amino acid residues 314-338 of  $\alpha$ -tubulin. The Examples also describe another embodiment (using the affinity probe referred to here as Probe 2) in  
10 which hemiasterlins bind to a region comprising about amino acid residues 200-300 of  $\alpha$ -tubulin, and more preferably comprising amino acid residues 204-280 of  $\alpha$ -tubulin.

It is understood, therefore, that hemiasterlins (including hemiasterlin derivatives and analogs such as HTI-286) can be expected to bind to either one or both of these regions on  $\alpha$ -tubulin. Moreover, mutations in either one or both of these regions are  
15 expected to disrupt the binding of such hemiasterlin compounds to tubulin. Consequently, cells having such mutations may be resistant to certain anti-tubulin compounds and, in particular, to hemiasterlin compounds (including hemiasterlin derivatives and analogs such as HTI-286). Hence, it is possible to identify cells (including cancer and tumor cells, such as ones obtained from a biopsy of a patient or  
20 other individual) that are either resistant to or are expected to be resistant to treatment with a hemiasterlin compound, by identifying cells that express tubulin having one or more mutations a binding regions identified with an affinity probe. Such methods are therefore encompassed by the present invention.

Generally speaking, to identify such resistant cells and tumors, it is necessary to  
25 determine the nucleotide or protein sequence of the tubulin (in particular,  $\alpha$ -tubulin) expressed by a cell or tumor. Generally, it will not be necessary to determine the complete amino acid or nucleotide sequence, but is sufficient to determine only the sequence of one or more binding sites identified by an affinity probe, such as one or more of the binding sites described *supra*. Using the methods described here it is possible to  
30 determine the relevant sequence of tubulin nucleic acid or protein in cells (*e.g.* derived from cell culture) or tissues from an individual, such as in cells or tissues in a sample obtained or derived from an individual subject or patient. These samples can be derived from, for example, a biopsy sample of a patient suspected of having a type of cancer or

other disorder associated with tubulin, or suspected of having a propensity for such a cancer or other disorder.

A variety of methods known in the art may be used to detect assay levels of mutated tubulin sequences in a sample. Diagnostic methods for the detection of tubulin nucleic acids in patient samples or in other cell or tissue sources may involve their  
5 amplification, *e.g.*, by PCR (see, for example, the experimental embodiment taught in U.S. Patent No. 4,683,202) followed by nucleotide sequencing of the amplified molecules using techniques that are well known to those of skilled in the art.

Preferably, the nucleotide or amino acid sequence obtained from a cell or tumor is  
10 compared to a consensus or wild-type sequence for tubulin, such as one of the  $\alpha$ -tubulin sequences depicted in **Figure 12**, or a suitable homolog or ortholog thereof to which an affinity probe of the invention binds or is expected to bind, or which is known to be susceptible to treatment with a hemiasterlin compound. In embodiments where the nucleotide sequence (such as a cDNA or mRNA sequence) encoding a tubulin molecule  
15 is determined, it is generally preferable to determine the tubulin amino acid sequence that nucleic acid is expected to encode, to determine whether one or more amino acid residues in a binding region is altered or mutated.

As explained above, the amino acid sequence of  $\alpha$ -tubulin is highly conserved among a number of different species of organisms and, in particular, among different  
20 species of vertebrates, as shown in **Figure 12**. Hence, the amino acid residues recited above and throughout this specification, unless otherwise noted, specifically refer to amino acid residues in the sequence of human  $\alpha$ -tubulin set forth in **Figure 12**. However, those skilled in the art will be able to readily identify corresponding domains and regions of other  $\alpha$ -tubulin amino acid sequences, including  $\alpha$ -tubulin sequences from other  
25 species of organisms, orthologs and homologs. For instance, residues of other  $\alpha$ -tubulin sequences shown in **Figure 12** that align with the amino acid sequences recited above are also understood to represent binding domains for hemiasterlin compounds.

Those skilled in the art can also readily align different  $\alpha$ -tubulin or other amino acid sequences, using routine algorithms such as FASTA (Pearson & Lipman, *Proc. Natl. Acad. U.S.A.* 1988, 85:2444-2448; Pearson, *Methods Enzymol.* 1990, 183:63-98),  
30 BLAST (Altschul *et al.*, *Nucl. Acids Res.* 1997, 25:3389-3402; Altschul, *J. Mol. Evol.* 1993, 36:290-300; Altschul *et al.*, *J. Mol. Biol.* 1990, 215:403-410), CLUSTAL and CLUSTALW (Higgins *et al.*, *Nucl. Acids. Res.* 1994, 22:4673-4680), to name a few.

Generally, such alignment algorithms will be used with the standard or default parameters, including standard alignment scoring systems and/or a scoring matrix such as BLOSUM62. See, Henikoff & Henikoff, *Proc. Natl. Acad. Sci.* 1992, 89:10915-10919. However, in certain circumstances that will be appreciated by those skilled in the art, it may be preferable to use nonstandard parameters and/or scoring matrices. For example, in 5 embodiments where very similar amino acid sequences are being compared (such as sequences of  $\alpha$ -tubulin) it may be preferable to use a scoring matrix such as BLOSUM90, that has higher cutoffs.

Preferred embodiments of such a detection scheme include the use of genomic 10 DNA or the synthesis of a cDNA molecule from an RNA molecule of interest (*e.g.*, by reverse transcription). A sequence within the DNA may then be used as a template for a nucleic acid amplification reaction such as PCR. Nucleic acid reagents used as synthesis initiation reagents (*e.g.*, primers) in the reverse transcription and amplification steps of such an assay are preferably chosen from the tubulin nucleic acid sequences or are 15 fragments thereof. Preferably, the nucleic acid reagents are at least about 9 to 30 nucleotides in length. PCR product can then be sequenced by standard methods to those familiar with the art.

## 7. EXAMPLES

20 The present invention is also described and demonstrated by way of the following examples. However, the use of these and other examples anywhere in the specification is illustrative only and in no way limits the scope and meaning of the invention or of any exemplified term. Likewise, the invention is not limited to any particular preferred embodiments described here. Indeed, many modifications and variations of the invention 25 may be apparent to those skilled in the art upon reading this specification, and such variations can be made without departing the invention in spirit or in scope. The invention is therefore to be limited only by the terms of the appended claims along with the full scope of equivalents to which those claims are entitled.

### 30 7.1. Materials and Methods

#### 7.1.1 Reagents

HTI-286 (N, $\beta$ -trimethyl- L -phenylalany- $N^1$ [(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]- $N^1$ ,3-dimethyl- L -valinamide trifluoroacetate, also known as SPA 110, was

synthesized based on methods reported previously (Nieman *et al.*, *J. Nat. Prod.* 2003, 66:183-199) that were subsequently modified (Zask *et al.*, *Proc. Am. Assoc. Cancer Res.* 2002, 43:737). Two benzophenone photoaffinity analogs of HTI-286 were made (Kaplan *et al.*, *National Medicinal Chemistry Symposium* 2002, abstract 49). The first analog, which is referred to here as Probe 1, is the compound 4-benzoyl-N, $\beta$ ,  $\beta$ -trimethyl-L-phenylalanyl-  $N^1$  -[(1 S, 2E)-3-carboxy-1-isopropylbut-2-enyl]- $N^1$ , 3-dimethyl-L-valinamide (Kaplan *et al.*, *National Medicinal Chemistry Symposium* 2002, abstract 49). The second analog, referred to here as Probe 2, is the compound N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1 S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N, $\beta$ , $\beta$ -trimethyl-L-phenylalaninamide. The chemical structure and other properties of these HTI analogs are summarized in **Table 1**, below.

Except where otherwise stated, experiments are done using microtubule associated protein (MAP)-rich bovine brain tubulin. MAP-rich bovine brain tubulin (99% purity), HeLa cell tubulin (90 % purity), guanosine-5'-triphosphate (GTP) and PEM buffer (80 mM Piperazine-N,N'-bis(2-ethanesulfonic acid) containing 1 mM ethylenebis(oxyethylenenitrilo) tetraacetic acid (EGTA) and, 1 mM magnesium chloride, pH = 6.8)) were obtained from Cytoskeleton (Denver,CO). Paclitaxel, vinblastine, colchicine, leupeptin, dimethylsulfoxide (DMSO), phenylmethanesulfonyl fluoride (PMSF), sodium dodecyl sulfate (SDS) and subtilisin were obtained from Sigma (St.Louis, MO). Dolastatin-10 was obtained from the National Cancer Institute. Probe 1, Probe 2, HTI-286, paclitaxel, vinblastine, colchicine, and dolastatin-10 were prepared as 1 or 10 mM stocks in DMSO for competition studies. Trypsin treated with L-(tosylamido-2-phenyl) ethyl chloromethyl ketone (TPCK) was obtained from Worthington Biochemical Corporation (Freehold, NJ). Formic acid, cyanogen bromide (CNBr), iodoacetamide (IAM), dithiothreitol (DTT), and trifluoroacetic acid (TFA) were obtained from Pierce (Rockford, IL). Guanidine hydrochloride was obtained from the Aldrich Chemical Co., Inc. (Milwaukee, WI). Acetic acid, methanol, and acetonitrile were from Fisher Scientific (Fair Lawn, NJ). Beckman tissue solubilizer-450 (BTS-450) and Ready Protein, a liquid scintillation cocktail, were obtained from Beckman Coulter Instruments, Inc. (Fullerton, CA). EN<sup>3</sup>HANCE autoradiography enhancer was obtained from NEN<sup>TM</sup> Life Science Products, Inc. (Boston, MA). Precast gels used for electrophoresis were obtained from BioRad (Hercules, CA).



KB cells are obtained from the American Type Culture Collection (Manassas, VA) accession no. CCL-17.

#### 7.1.2 Reversible binding affinity assays

5       The affinity of both HTI-286 and of non-radiolabeled Probe 1 and Probe 2, to bovine brain tubulin was determined by previously published methods (Krishnamurthy *et al.*, *Biochem.*, submitted). Briefly, 5 mg/mL rhodamine-labeled tubulin prepared in PEM buffer was pre-incubated with buffer or varying concentrations of test agents in individual quartz cuvettes. The intrinsic tryptophan fluorescence of tubulin (excitation at 560 nm  
10       and maximal emission at 580 nm) was measured after pre-incubation for 90 minutes. The change in fluorescence intensity at 580 nm in the presence of compound was fitted to a quadratic equation to obtain apparent dissociation constants ( $K_D$ ) of the test agents.

      Binding of Probe 1 to bovine brain tubulin was also assessed by incubating 0.5  $\mu$ M tubulin with 0.005-5  $\mu$ M radiolabeled Probe 1 both in the presence and in the absence  
15       of 5  $\mu$ M non-radiolabeled probe 1 as a competitor. Incubations were performed for 30 minutes at room temperature in PEM buffer. Binding was measured using centrifugal gel filtration on Sephadex G-50 (Microspin G-50) as described (Bai *et al.*, *Cancer Res.* 1996, 56:4398-4406). Spin columns (Amersham Biosciences, Piscataway, NJ) were centrifugated for 1 minute at 2000 rpm in a microcentrifuge immediately prior to use, and  
20       samples applied in 50  $\mu$ L total volume, followed by a 2 minute centrifugation at the same speed. The amount of probe 1 bound to tubulin was quantitated by measuring the radioactivity within aliquots collected after centrifugation. The  $K_D$  value, representing 50% of specific binding, was calculated from the curve generated by subtracting the amount of radiolabeled probe bound in the absence of probe from that bound in the  
25       presence of non-radiolabeled probe.

#### 7.1.3 Tubulin polymerization assay in a cell-free system

*In vitro* tubulin polymerization assays were performed according to procedures that have been described elsewhere (Loganzo *et al.*, *Cancer Res.* 2003, 63:1838-1845).  
30       Briefly, bovine MAP-rich tubulin (final concentration 1.5 mg/mL) was dissolved in cold PEM buffer containing 1 mM GTP (GPEM) and centrifuged at 12,000 x g for 10 minutes at 4°C. The tubulin solution (100  $\mu$ L/well) was added rapidly to wells of a low-volume, 96-well plate already containing duplicate aliquots (10  $\mu$ L) of test compounds in GPEM. Final compound concentrations were 0.3  $\mu$ M. Control wells contained the same final

concentration of DMSO (0.3%). After initiation of the reaction, absorbance at 340 nm was measured every minute for 60 minutes at 24°C using a SpectraMax Plus plate reader (Molecular Devices, Sunnyvale, CA). An increase in absorbance over time indicated an increase in turbidity resulting from tubulin polymerization.

5

#### 7.1.4 Cell proliferation assay in tissue culture

Cell proliferation studies were performed according to procedures described by Loganzo *et al.* (*Cancer Res.* 2003, 63:1838-1845). Briefly, KB cells were plated in 96-well plates in 100 µL media at densities pre-determined to produce 60-90% confluence at the time of analysis. Compounds, which were serially diluted into media as 2x stocks, were added to cells in duplicate. After 72 hours of incubation, cell survival was assayed by the SRB assay as described (Rabindran *et al.*, *C. Cancer Res.* 1998, 58:5850-5858). IC<sub>50</sub> values (*i.e.* the concentration of drug needed to inhibit cell growth by 50%) were used to evaluate drug potency.

15

#### 7.1.5 Photolabeling, SDS-PAGE and fluorography techniques

Photoaffinity labeling studies were done by incubating tubulin (2.5 µM and 5 µM for Probe 1 and 2 experiments, respectively, unless otherwise noted) prepared in PEM buffer with or without competitor molecule for 15 minutes at 4°C, room temperature, or 37°C, prior to incubation with radiolabeled probe (2.5 or 0.25 µM) at room temperature or 37°C for 30 minutes. Alternatively, a cytosolic preparation of KB-3-1 cell lysates (50 µg) or KB-3-1 cells (40,000 cells) were used in the place of purified tubulin above. The supernatant of mechanically disrupted cells centrifuged at 100,000 x g was used as a cytosolic preparation. K-3-1 cells were prepared by growing KB-3-1 cells in a 96-well plate and rinsing them with serum free medium. For experiments with whole cells, 250 µM HTI-286 was used as the competitor molecule during pre-incubation at 37°C for 15 minutes.

Samples were irradiated at 360 nm with a Mineralight lamp (UVP, Upland, CA) for 30 minutes (for Probe 1), 2 hours (for whole cells with Probe 1), or 2 hours (for Probe 2) at 4°C, unless otherwise noted. Radiation of the benzophenone moiety at this wavelength effectively activates the photoprobe with minimal destruction of protein compared to the shorter wavelengths typically used for azido-containing photoprobes

30

(see, Williams *et al.*, *Methods Enzymol.* 1986, 126:667-682). If whole cells were used, Laemmli sample buffer was then added to lyse the cells.

Photolabeled samples were analyzed by SDS-PAGE using 7.5% Tris-HCl gels. Low grade SDS (Sigma, catalog No. L5750) was used in the running buffer, when  
5 needed, to allow separation of  $\alpha$  and  $\beta$ -tubulin subunits. Gels were fixed, stained with Coomassie blue from BioRad (Hercules, CA) to ensure equal loading of protein, and sliced. The radioactivity of each 1 mm slice was determined by placing the slice overnight at room temperature in 200  $\mu$ L of 90% BTS-450, followed by the addition of 6 mL Ecolume scintillation cocktail (for Probe 1) or Ready Protein scintillation cocktail  
10 (for Probe 2) before subjecting the vials to scintillation counting. Alternatively, gels were incubated with EN<sup>3</sup>HANCE according to the manufacturer's instructions (Life Sciences Products, Inc.), washed, and dried prior to exposure of the gel to Biomax MX film from Kodak (Rochester, NY) for fluorography.

#### 15                    7.1.6 Formic Acid digestion after photoaffinity labeling using Probe 1

50 $\mu$ M tubulin, which was photolabeled with 5  $\mu$ M (7.2  $\mu$ Ci) [<sup>3</sup>H] Probe 1 according to methods *supra*, was run on a 7.5% Tris-HCl SDS-PAGE gel that allowed the  $\alpha$  and  $\beta$ -tubulin to co-migrate. Tubulin was excised from the gel by cutting the unstained  
20 gel in the region that co-migrated precisely with the 50 kDa marker of the Precision Protein Standards (BioRad). The material was digested in-gel in 250  $\mu$ L of 75% formic acid at 37°C. After 44-72 hours, formic acid was removed in a speed vacuum, and digestion products were separated by SDS-PAGE on Tris-tricine gels (10-20%). Radiolabeled peptides were visualized by fluorography. In other types of experiment, tubulin subunits were separated on 7.5% Tris-HCl gels (BioRad) and  $\alpha$ - and  $\beta$ -tubulin  
25 bands were cut and digested separately. The sequence of the labeled formic acid digestion fragment was confirmed by mass spectrometric (MS) analysis as described below.

#### 7.1.7 Trypsin digestion of native tubulin after photoaffinity labeling

Native tubulin (15  $\mu$ g for Probe 1 or 5  $\mu$ g for Probe 2) was incubated with 0.25  
30  $\mu$ M (0.35  $\mu$ Ci) of [<sup>3</sup>H]-Probe 1 or 2.5  $\mu$ M [<sup>3</sup>H]-probe 2 in PEM buffer at room temperature for 30 minutes. Samples were then irradiated at 4°C for 2 hours and digested with 0.8  $\mu$ g trypsin for 5 to 60 minutes at 30°C. The reaction was stopped by adding leupeptin to 0.01 mM. The samples were separated on 10-20% Tris-tricine gels. The

radioactivity within 1 mm gel slices was determined. Alternatively, peptides were resolved on 7.5% Tris-HCl and gels were exposed to film using fluorographic methods described above.

5                    7.1.8 CNBr cleavage after Photoaffinity Labeling of Tubulin

Tubulin (2.5  $\mu$ M) was photolabeled with 0.25  $\mu$ M [ $^3$ H]-Probe 1 or 2.5  $\mu$ M [ $^3$ H]-Probe 2 as described above.  $\alpha$ - and  $\beta$ -isomers were then resolved on a 7.5% Tris-Glycine SDS gel for Probe 1 or a 7.5% Tris-HCl SDS gel for Probe 2. Gel bands were independently excised, reduced with dithiothreitol (10 mM in 100mM ammonium bicarbonate) and alkylated with iodoacetamide (25 mM in 100 mM ammonium bicarbonate).

The gel bands containing separated tubulins were shrunk with 100% acetonitrile and reswelled with CNBr (150 mg/ml) in 70% formic acid. Digestion was allowed to proceed for 4.5 hours at room temperature. Gel bands were then evaporated to dryness from H<sub>2</sub>O several times in vacuum to remove excess reagents. The gel bands were then reswollen with SDS loading buffer and mounted on top of a 10-20% Tris-tricine gel where digestion products were separated. Replicate gels were run of the CNBr fragments: one gel was electroblotted to PVDF for autoradiography and the other was silver stained (Shevchenko *et al.*, *Analytical Chemistry* 1996, 68:850-858). Blots for autoradiography were first soaked with a solution of PPO in toluene and autoradiographic images were captured on Kodak BioMax MS film.

As in the formic acid digestion protocol (see Section 7.1.6), the sequence of the labeled CNBr digestion fragment was confirmed by mass spectrometric analysis from the silver stained gel as described below.

25

7.1.9 Mass Spectrometric Analyses of Digestion Fragments

Peptides from the in-gel enzymatic digestion were injected onto a self-packed PicoFrit C18 column from New Objectives (Woburn, MA) that is directly interfaced to an LCQ Deca ion trap mass spectrometer from ThermoFinnigan (San Jose, CA). During a 90 minute HPLC gradient [4% to 60% solvent B (0.1 M acetic acid/90% ACN/H<sub>2</sub>O), solvent A: 0.1 M acetic acid/H<sub>2</sub>O], the mass spectrometer was operated in a data-dependent mode using software provided by the manufacturer to acquire both MS (peptide mass) and MS/MS (fragment ion mass) spectra. Peptide sequences were

determined by matching the fragment ion spectra against the non-redundant NCBI protein database using the Sequest search algorithm provided by ThermoFinnigan.

#### 7.1.10 Subtilisin Digestion of Tubulin before or after Labeling with Probe 1

5        Before or after incubating with 0.25  $\mu$ M (0.36  $\mu$ Ci) [ $^3$ H]-Probe 1 for 30 minutes at room temperature and irradiating as described in Section 7.1.5, tubulin (5  $\mu$ g) was digested under native conditions with subtilisin (0.2  $\mu$ g) for 60 minutes at 30°C. Enzymatic digestions were stopped with 1 mM PMSF. After both digestion and labeling, samples were resolved by SDS-PAGE under conditions that allowed separation of  $\alpha$ - and  
10     $\beta$ -tubulin subunits. Gels were then fixed and stained with Coomassie blue from BioRad (Hercules, CA) and subjected to fluorography.

#### 7.1.11 Formic Acid and Lys C Digestion of Tubulin after Labeling with Probe 2

15        Probe 2 photolabeled tubulin (as described in Section 7.1.5) was digested with formic acid (75%) or Lys C (0.1  $\mu$ g) in the presence of 0.1% SDS at 37°C for 72 hours. The samples were separated on 10-20 % Tris-tricine gels. The gels were exposed to film using fluorographic methods described above.

#### 7.1.12 Subtilisin Digestion of Tubulin before Labeling with Probe 2

20        Tubulin was digested with 0.2  $\mu$ g of subtilisin for 60 min at 30°C and the reaction was stopped by the addition of PMSF (1 mM). Digested tubulin was then photolabeled with Probe 2 as described in Section 7.1.5. The samples were separated on 10-20% Tris-tricine gels. The gels were exposed to film using fluorographic methods described in  
25    Section 7.1.5.

#### 7.1.13 Subtilisin Digestion of Tubulin after Incubation of Tubulin with Various Compounds

30        Tubulin (5  $\mu$ g) was incubated with 50  $\mu$ M drug at room temperature for 25 minutes and then digested under native conditions with subtilisin (0.05  $\mu$ g) for 0, 5, 10, 15, 30 and 60 min at room temperature. Enzymatic digestion was stopped with 1 mM PMSF and samples were resolved by SDS-PAGE under conditions that allowed separation of  $\alpha$ - and  $\beta$ -tubulin subunits. Gels were then fixed and stained with Coomassie blue.

#### 7.1.14 Separation of Polymerized and Free Tubulin

Tubulin (5  $\mu$ g) was incubated with 50  $\mu$ M drug at room temperature for 25 minutes and then centrifuged at either 10,000 g for 15 minutes in a tabletop microfuge or at 100,000 g for 15 minutes in an airfuge. The supernatant was carefully removed to a fresh tube to obtain free tubulin and the pellet was solubilized in SDS-PAGE sample buffer. The free and polymerized tubulin were run on a 7.5% Tris-HCl gel and gels were stained with Coomassie blue.


#### 10           7.2.   Synthesis of Hemiasterlin Analog Probes

Probes were designed to mimic the binding of hemiasterlin and its analogs. SARs (structure activity relationship) studies on hemiasterlins and their analogs (Nieman *et al.*, *J. Nat. Prod.* 2003, 66:183-199; Zask *et al.*, *Proc Am. Assoc. Cancer Res.* 2002, 43:737) have determined which structural modifications in hemiasterlin allow the molecule to retain activity. Probe designs were constrained to only these modifications. More than one probe was designed and synthesized with the intent to define the binding at more than one point of contact. The syntheses and radiolabeling of Probe 1 and Probe 2 are described in the sections, *infra*.

#### 20           7.2.1 Synthesis of Probe 1 and Probe 2 precursors

**Scheme I**, below, diagrams the synthesis of N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-4-(2-phenyl-1,3-dioxolan-2-yl)-L-phenylalanine (Compound 9) and 4-benzoyl-N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-L-phenylalanine (Compound 8) that are precursors for Probe 1 and Probe 2, respectively.


25

**Scheme I**

The syntheses, below, describe the preparation of the compounds shown in


5 **Scheme I.**

Synthesis of  
methyl 3-methyl-3-[(4-benzoyl)phenyl]butanoate (Compound 3)



According to Scheme I, a round-bottomed flask is charged with methyl 3-methyl 3-phenyl butanoate (0.25 g, 1.3 mmol, Compound 1), benzoyl chloride (0.15 mL, 1.3 mmol, Compound 2), and carbon disulfide (1.6 mL). While stirring under nitrogen atmosphere, the reaction mixture is cooled to 0 °C in an ice-water bath. Aluminum chloride (0.35 g, 2.6 mmol) is added in a single portion and the cooling bath is removed. The reaction mixture is heated at reflux 4 hours and then allowed to cool to room temperature. The reaction mixture is transferred dropwise into ice-water. The aqueous phase is extracted thrice with dichloromethane. The combined organic extracts are washed with water and 5 % aqueous potassium carbonate, dried over sodium sulfate, decanted, and concentrated under reduced pressure to give a brown liquid. MS (ES<sup>+</sup>): m/z (M+H) = 297.5

Synthesis of  
methyl 3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoate (Compound 4)




According to Scheme I to a solution of methyl 3-methyl-3-[(4-benzoyl)phenyl]butanoate (26 mmol max, Compound 3) in toluene (100 mL), ethylene glycol (3.2 g, 52 mmol) and *p*-toluenesulfonic acid monohydrate (~ 10 mg) are added. After heating at reflux for 2 hours, an additional 5 mL ethylene glycol and a Dean-Stark trap are added, and reflux is re-started. Azeotropic distillation of water is allowed to proceed overnight. A mixture of ethylene glycol and water (8 mL) is observed in the Dean-Stark trap. The LC/MS of an aliquot of the reaction mixture revealed the presence of both starting material and desired product. The reaction mixture is concentrated under reduced pressure and the residue is partitioned between diethyl ether and saturated aqueous sodium hydrogen carbonate. The aqueous phase is extracted thrice with diethyl ether. The combined extracts are washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride, dried over sodium sulfate, decanted, and concentrated under reduced pressure. The crude product is taken up in toluene (100



mL); and ethylene glycol (3.2 g, 52 mmol) and *p*-toluenesulfonic acid monohydrate (~ 10 mg) are added. The reaction mixture is heated at reflux for 4 hours without the Dean-Stark apparatus and then overnight with the trap. After cooling the reaction mixture to room temperature, the aqueous work-up above is performed. The crude residue is  
5 purified by flash chromatography (ethyl acetate/hexanes) to furnish (2.6 g, 29 % for 2 steps) of an amorphous white solid. TOF MS ( $ES^+$ ): ( $M+H$ ) = 341.3

## Synthesis of

3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoic acid (Compound 5)


10

According to Scheme I to a suspension of methyl 3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoate (2.6 g, 7.6 mmol, Compound 4) in tetrahydrofuran (20 mL), methanol (20 mL), and water (10 mL), lithium hydroxide monohydrate (0.48 g, 11  
15 mmol) is added. The mixture is heated at 55 °C for 5 hours, during which all solids dissolved. The reaction mixture is then allowed to cool to room temperature and solvents are evaporated under reduced pressure. The white solid is partitioned between ethyl acetate and water. Most of the material remained undissolved. The biphasic mixture is cooled to 0 °C in an ice-water bath. Glacial acetic acid is added in portions until pH = 5.  
20 At this point, white solid precipitated, leaving a clear, colorless supernatant. Ethyl acetate is then removed under reduced pressure, and the solids are isolated by filtration of the aqueous phase and washed with cold water. After drying, a white solid is obtained (2.3 g, 92 %). TOF MS ( $ES^+$ ): ( $M+H$ ) = 327.2

## Synthesis of

(4S)-3-{{(2S)-2-azido-3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoyl}-4-benzyl-1,3-oxazolidin-2-one (Compound 6a) and (4S)-3-{{(2S)-2-azido-3-(4-benzoylphenyl)-3-methylbutanoyl}-4-benzyl-1,3-oxazolidin-2-one (Compound

5

6)

According to Scheme I to a solution of 3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoic acid (2.3 g, 7.0 mmol, Compound 5) in anhydrous tetrahydrofuran (14 mL) under a nitrogen atmosphere, triethylamine (1.2 mL) is added. The mixture is cooled to -78°C in a dry-ice acetone bath. Pivaloyl chloride (0.91 mL, 7.4 mmol) is added dropwise, causing the immediate formation of a white precipitate. The reaction mixture is allowed to sit for 20 minutes at -78°C and is then stirred at 0 °C in an ice-water bath. In a separate flask, a solution of *S*-benzyloxazolidinone (1.2 g, 6.9 mmol) in anhydrous tetrahydrofuran is prepared under a nitrogen atmosphere and cooled to -35°C in a dry-ice/acetone bath. A small amount of triphenylmethane (< 5 mg) is added as an indicator of deprotonation. *n*-butyllithium (1.6 M solution in hexanes, 4.5 mL, 7.2 mmol) is added dropwise via syringe. At the end of this addition, the characteristic pinkish orange color of the anion of triphenylmethane is not yet observed. However, after the introduction of an additional 0.2 mL of *n*-butyllithium, this color is achieved. After 30 minutes of stirring at 0 °C, the flask containing the mixed anhydride is re-cooled to -78°C in a dry-ice/acetone bath. The solution of the lithium anion of the oxazolidinone is added to the mixed anhydride solution via a cannula. The source flask is washed twice with tetrahydrofuran (4 mL × 2) and these washings are also transferred via a cannula to the mixed anhydride solution. The reaction mixture is stirred at -78°C for 1 hour and at 0 °C for 1 hour, and then allowed to warm to room temperature overnight. Water (~ 15 mL) is added and stirring is continued for 10 minutes. The aqueous phase is extracted thrice with diethyl ether. The combined extracts are washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride, dried over sodium sulfate,

decanted, and concentrated under reduced pressure to afford (3.6 g, > 100 % crude) of a white foam. MS (ES<sup>+</sup>): m/z (M+H) = 486.2


A solution of this crude benzophenone ketal oxazolidinone (7.0 mmol maximum) in anhydrous tetrahydrofuran (40 mL) is cooled to -78 °C in a dry-ice/acetone bath while stirring under a nitrogen atmosphere. Potassium hexamethylsilazide (0.5 M solution in toluene, 18 mL, 9.0 mmol) is added dropwise to the solution via a syringe. A deep orange-red color resulted from this addition. After stirring for 1 hour at -78 °C, a pre-cooled solution of triisopropylsulfonyl azide (3.0 g, 9.8 mmol) in tetrahydrofuran (20 mL) at the same temperature is added rapidly via a cannula. After stirring for 3 minutes at -78 °C, the reaction mixture is quenched by the addition of glacial acetic acid (1.8 mL), which caused a color change from deep red to pale yellow. The cooling bath is removed and the reaction is stirred at room temperature for 20 minutes, followed by 1 hour at 40 °C. After cooling to room temperature, the reaction mixture is diluted with water and extracted thrice with diethyl ether. The combined extracts are washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride, dried over sodium sulfate, decanted, and concentrated under reduced pressure to afford a pale yellow oil. This oil is inert to hydrolysis of the ketal by two methods: treatment with *p*-toluenesulfonic acid in aqueous acetone and with aqueous hydrochloric acid in tetrahydrofuran. The unaffected crude material is purified by flash chromatography (hexanes/ethyl acetate) to afford a clean separation of (4S)-3-((2S)-2-azido-3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoyl]-4-benzyl-1,3-oxazolidin-2-one (0.75 g, 1.4 mmol, Compound 6a) and (4S)-3-((2S)-2-azido-3-(4-benzoylphenyl)-3-methylbutanoyl]-4-benzyl-1,3-oxazolidin-2-one (0.82 g, 1.7 mmol, Compound 6), giving a total yield (3.1 mmol / 7.0 mmol) of 44 % for three steps -- the formation of the mixed anhydride, displacement with the lithium oxazolidinone, and preparation of the azide: (4S)-3-((2S)-2-azido-3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoyl]-4-benzyl-1,3-oxazolidin-2-one: TOF MS (ES<sup>+</sup>) = 527.4 (4S)-3-((2S)-2-azido-3-(4-benzoylphenyl)-3-methylbutanoyl]-4-benzyl-1,3-oxazolidin-2-one: TOF MS m/z (ES<sup>+</sup>) = 483.4

30

#### Synthesis of

(alphaS)-4-Benzoyl-N-((4S)-3-[4-benzoyl-N-(tert-butoxycarbonyl)-β,β-dimethyl-L-phenylalanyl]-4-benzyl-2-oxo-1,3-oxazolidin-2-yl)-N-((4S)-3-[4-benzoyl-N-

(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-L-phenylalanyl]-4-benzyl-2-oxo-1,3-oxazolidin-4-yl}-N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-L-phenylalaninamide (Compound 7)



5 According to Scheme I, a solution of (4S)-3-{(2S)-2-azido-3-(4-benzoylphenyl)-3-methylbutanoyl]-4-benzyl-1,3-oxazolidin-2-one (0.80 g, 1.7 mmol, Compound 6) in ethyl acetate (8 mL) is degassed with a small piece of dry ice. When all effervescence had subsided, palladium on carbon (Pd/C, 10 %, 10 mg) is added in a single portion, followed by di-*t*-butyl dicarbonate (0.74 g, 3.4 mmol). The reaction flask is evacuated  
10 under weak house vacuum and then flushed with hydrogen (balloon pressure). This process is repeated thrice. Finally, the reaction mixture is allowed to stir under a hydrogen atmosphere. After 30 minutes, the reaction is incomplete according to thin-layer chromatography (TLC, 20 % ethyl acetate/hexanes); hence, stirring under hydrogen is continued over the weekend (~ 64 hours). Following this interval, TLC showed a  
15 complete disappearance of starting material and the emergence of two new spots. LC/MS analysis revealed these products to be both the desired material and the benzyl phenyl alcohol, the by-product of ketone reduction. The reaction mixture is filtered through a Diatomaceous earth pad to remove Pd/C. The filtrate is concentrated under reduced pressure to afford a clear, colorless oil. This material is subjected to manganese (IV)  
20 oxide in dichloromethane in order to oxidize the alcohol by-product back to the benzophenone. This method, however, proved to be very sluggish was aborted. The crude mixture (1.7 mmol max. of alcohol) was then taken up in dichloromethane (10 mL). Pyridinium dichromate (0.96 g, 2.6 mmol) was added to the solution and the resulting rust-colored mixture is stirred overnight at room temperature. TLC showed a complete  
25 conversion of by-product to desired product. The reaction mixture is filtered through a Diatomaceous earth pad to remove most of the chromium salts. The filtrate is concentrated under reduced pressure to a dark brown oil, and this crude material is purified by flash chromatography (ethyl acetate/hexanes) to afford (0.52 g, 55 %) of a


hard, white foam. An additional 0.24 g of slightly impure material is also recovered and removed.

TOF MS  $m/z$  ( $ES^+$ ) = 557.5

5


### Synthesis of

4-benzoyl-N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-L-phenylalanine (Compound 8)



According to Scheme I to a 0 °C solution of (alphaS)-4-benzoyl-N-((4S)-3-[4-benzoyl-N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-L-phenylalanyl]-4-benzyl-2-oxo-1,3-oxazolidin-2-yl)-N-((4S)-3-[4-benzoyl-N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-L-phenylalanyl]-4-benzyl-2-oxo-1,3-oxazolidin-4-yl)-N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-L-phenylalaninamide (0.49 g, 0.88 mmol, Compound 7) in tetrahydrofuran (11 mL) and water (3 mL), hydrogen peroxide (30 % aqueous solution, 0.76 mL, 7.9 mmol) followed by lithium hydroxide monohydrate (0.11 g, 2.6 mmol) are added. The reaction mixture is stirred for 23 hours while gradually warming to room temperature. The reaction is quenched by the addition of sodium sulfite (1.5 M aqueous solution, 10 mL, 15 mmol), which is accompanied by slight exothermicity. The quenched mixture is stirred for 1 hour at room temperature and then cooled to 0 °C in an ice-water bath. The pH of the mixture is adjusted to 4 by the addition of citric acid (1 M aqueous solution). The acidified mixture is then extracted thrice with ethyl acetate. The combined extracts are washed with saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, decanted, and concentrated under reduced pressure to afford a white foam (0.47 g). This crude product is dissolved in acetonitrile/water (1:1) and purified by semi-preparative reverse-phase HPLC, employing a gradient elution of 5 % acetonitrile/95 % water to 100 % acetonitrile over 1 hour. A hard, white foam (0.23 g, 66 %) is obtained after collection and concentration. TOF MS ( $ES^-$ ):  $m/z$  (M-H) = 396.2

Synthesis of  
N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-4-(2-phenyl-1,3-dioxolan-2-yl)-L-phenylalanine (Compound 9)



5

According to Scheme I, a solution of (4S)-3-((2S)-2-azido-3-methyl-3-[4-(2-phenyl-1,3-dioxolan-2-yl)phenyl]butanoyl}-4-benzyl-1,3-oxazolidin-2-one (0.74 g, 1.4 mmol, Compound 6a) in ethyl acetate (25 mL) is degassed with a small piece of dry ice. When all effervescence had subsided, palladium on carbon (Pd/C, 10 %, 20 mg) is added in a single portion, followed by di-*t*-butyl dicarbonate (0.61 g, 2.8 mmol). The reaction flask is evacuated under weak house vacuum and then flushed with hydrogen (balloon pressure). This process is repeated thrice. Finally, the reaction mixture is allowed to stir under a hydrogen atmosphere. After 4 hours, TLC showed a complete disappearance of starting material and the emergence of a single new spot of lower retention factor. The reaction mixture is filtered through a Diatomaceous earth pad to remove Pd/C. The filtrate is concentrated under reduced pressure to afford (1.3 g, >100 %) of a clear, light blond oil.


To a 0 °C solution of this crude material (1.3 g, 1.4 mmol maximum) in tetrahydrofuran (17 mL) and water (4 mL), hydrogen peroxide (30 % aqueous solution, 1.4 mL, 13 mmol) followed by lithium hydroxide monohydrate (0.18 g, 4.2 mmol) are added. The reaction mixture is stirred for 23 hours at room temperature. LC/MS analysis revealed the reaction to be incomplete. The mixture is cooled to 0 °C and an additional 4 mL of hydrogen peroxide solution and 0.18 g of lithium hydroxide monohydrate is added. Stirring is continued for 60 hours while the mixture gradually warmed to room temperature. The reaction is quenched by the addition of sodium sulfite (1.5 M aqueous solution, 25 mL, 38 mmol), which is accompanied by slight exothermicity. The quenched mixture is stirred for 1 hour at room temperature and then cooled to 0 °C in an ice-water bath. The pH of the mixture is adjusted to 4 by the addition of citric acid (1 M aqueous solution). The acidified mixture is then extracted thrice with ethyl acetate. The combined

extracts are washed with a saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, decanted, and concentrated under reduced pressure to afford a white foam (0.90 g). This crude product is dissolved in dimethylsulfoxide and purified by semi-preparative reverse-phase HPLC, employing a gradient elution of 5 %

- 5 acetonitrile/95 % water to 100 % acetonitrile over 1 hour. N-(tert-butoxycarbonyl)- $\beta,\beta$ -dimethyl-4-(2-phenyl-1,3-dioxolan-2-yl)-L-phenylalanine is obtained as a white powder (0.37 g, 60 %) after collection and concentration. TOF MS (ES<sup>-</sup>): (M-H) = 440.1

### 7.2.2 Synthesis of Probe 1


- 10 **Scheme II**, below, diagrams the synthesis of 4-benzoyl-N,  $\beta,\beta$ -trimethyl-L-phenylalanyl-N<sup>1</sup>-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N<sup>1</sup>,3-dimethyl-L-valinamide (Probe 1).



- 15 The syntheses, below, describe the preparation of the compounds shown in **Scheme II**.

## Synthesis of

N-(tert-butoxycarbonyl)-N,  $\beta,\beta$ -trimethyl-4-(2-phenyl-1,3-dioxolan-2-yl)-L-phenylalanine (Compound 10)



5


According to Scheme II, a solution of N-(tert-butoxycarbonyl)-  $\beta,\beta$ -dimethyl-4-(2-phenyl-1,3-dioxolan-2-yl)-L-phenylalanine (0.34 g, 0.77 mmol, Compound 9) in anhydrous dimethylformamide (11 mL) is cooled to 0 °C in an ice-water bath under a nitrogen atmosphere. Sodium hydride (60 % dispersion in mineral oil, 0.15 g, 3.9 mmol) is added slowly. After effervescence has ceased, methyl iodide (0.49 g, 7.8 mmol) is added via syringe. The reaction mixture is then allowed to warm to room temperature gradually while stirring overnight. The mixture is then cooled to 0 °C in an ice-water bath. Glacial acetic acid (1 mL) is added to adjust the pH to 4. The reaction mixture is partitioned between ethyl acetate and water. The aqueous phase is extracted thrice with diethyl ether. The combined extracts are washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride, dried over sodium sulfate, decanted, and concentrated under reduced pressure to a blond oil (0.49 g, > 100 %). MS (ES<sup>+</sup>): m/z (M+Na) = 492.3

The crude blond oil (0.49 g, 0.77 mmol maximum) is taken up in tetrahydrofuran (2 mL), methanol (2 mL), and water (1 mL). To this solution, lithium hydroxide monohydrate (81 mg, 1.9 mmol) is added. The reaction mixture is stirred for 24 hours at room temperature and an additional quantity of lithium hydroxide monohydrate (20 mg, 0.48 mmol) is added. Stirring is resumed for 60 hours, following which the solvent is evaporated under reduced pressure to give a white solid (0.62 g), which is purified by semi-preparative reverse-phase HPLC, employing a gradient elution of 5 % acetonitrile/95 % water to 100 % acetonitrile over 1 hour. N-(tert-butoxycarbonyl)-N,  $\beta,\beta$ -trimethyl-4-(2-phenyl-1,3-dioxolan-2-yl)-L-phenylalanine is obtained as a hard white foam (0.26 g, 74 % over 2 steps) after collection and concentration. TOF MS (ES<sup>-</sup>): m/z (M-H) = 454.1



## Synthesis of

4-benzoyl-N,  $\beta,\beta$ -trimethyl-L-phenylalanyl-N<sup>1</sup>-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N<sup>1</sup>,3-dimethyl-L-valinamide (Probe 1)



5

According to Scheme II, to a solution of N-(tert-butoxycarbonyl)-N,  $\beta,\beta$ -trimethyl-4-(2-phenyl-1,3-dioxolan-2-yl)-L-phenylalanine (0.23 g, 0.50 mmol, Compound 10) and ethyl (2E,4S)-2,5-dimethyl-4-[methyl(3-methyl-L-valyl)amino]hex-2-enoate (0.24 g, 0.76 mmol, Compound 11) in anhydrous dimethylformamide (3 mL) under a nitrogen atmosphere, hydroxybenzotriazole (0.14 g, 1.0 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.19 g, 1.0 mmol), and N-methylmorpholine (0.11 mL, 1.0 mmol) are added. After 24 hours, the mixture is diluted with water, and the aqueous layer is extracted with diethyl ether (3 times). The combined  
10 extracts are washed with 2 % hydrochloric acid and saturated aqueous sodium chloride, dried over sodium sulfate, filtered, and concentrated *in vacuo*. The residue is isolated as a hard white foam (0.34 g, 92 %, Compound 12). MS (ES<sup>+</sup>): m/z (M-Boc+H) = 650.4

To a 0 °C solution of this material in tetrahydrofuran (10 mL), 10 % hydrochloric acid (2.5 mL) is added. The mixture is stirred for 22 hours at room temperature then  
20 heated for 28 hours at 45 - 50 °C. After an additional 48 hours at room temperature, the reaction mixture is carefully quenched by the addition of saturated aqueous sodium hydrogen carbonate and then extracted thrice with diethyl ether. The combined organic extracts are washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride, dried over sodium sulfate, decanted, and concentrated under  
25 reduced pressure to afford a hard white foam (0.29 g, 91 %). MS (ES<sup>+</sup>): m/z (M-Boc+H) = 606.4


A solution of this material (0.29 g, 0.42 mmol) in dichloromethane (4 mL) is cooled to 0 °C in an ice-water bath. Trifluoroacetic acid (0.32 mL, 4.2 mmol) is added and the mixture is stirred for 30 minutes at 0 °C. The cooling bath is then removed.  
30 After an additional 2 hours of stirring, an additional quantity of trifluoroacetic acid (0.20

mL) is added. After 18 hours of stirring, solvent and excess acid are removed under reduced pressure to yield 0.43 g of a hard brown foam.

This material is taken up in tetrahydrofuran (2 mL), methanol (2 mL), and water (1 mL). Lithium hydroxide monohydrate (0.16 mg, 3.8 mmol) is added and the reaction mixture is stirred overnight at room temperature. The solvent is evaporated under reduced pressure. The residue is purified by semi-preparative reverse-phase HPLC (employing a gradient elution of 5 % acetonitrile/95 % water/0.1 % trifluoroacetic acid to 100 % acetonitrile over 1 hour) to give a hard, white foam (0.20 g). Subsequent isocratic reverse phase HPLC purifications (employing 60 % methanol/40 % water (0.02 % trifluoroacetic acid) furnished 4-benzoyl-N,  $\beta,\beta$ -trimethyl-L-phenylalanyl-N<sup>1</sup>-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N<sup>1</sup>,3-dimethyl-L-valinamide trifluoroacetic acid as a white powder (60 mg, 19 %). TOF MS (ES<sup>+</sup>): m/z (M+H) = 578.4

### 7.2.3 Synthesis of Probe 2


**Scheme III**, below, diagrams the synthesis of N, $\beta,\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N, $\beta,\beta$ -trimethyl-L-phenylalaninamide (Probe 2).



The syntheses, below, describe the preparation of the compounds shown in Scheme III.

### Synthesis of

N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N, $\beta$ , $\beta$ -trimethyl-L-phenylalaninamide (Probe 2)



According to Scheme III, to a solution of 4-benzoyl-N-(tert-butoxycarbonyl)- $\beta$ , $\beta$ -dimethyl-L-phenylalanine (0.13 g, 0.33 mmol, Compound 8), benzotriazole-1-yl-oxy-tris-pyrrolidinophosphonium hexafluorophosphate (0.26 g, 0.50 mmol), and dimethylaminopyridine (DMAP, 24 mg, 0.20 mmol) in dichloromethane (4 mL, Aldrich), diisopropylethylamine (0.17 mL, 0.99 mmol) is added under a nitrogen atmosphere. To this mixture is added a solution of ethyl (E,4S)-2,5-dimethyl-4-(methylamino)-2-hexenoate (0.38 g, 1.1 mmol, Compound 13) in anhydrous dichloromethane (3 mL). The resulting reaction mixture is stirred at room temperature for 18 hours. Volatiles are evaporated under reduced pressure. The crude product (0.70 g) is purified by semi-preparative HPLC (employing a gradient elution of 5 % acetonitrile/95 % water to 100 % acetonitrile over 1 hour). A hard, white foam (0.16 g, 84 %, Compound 14) is obtained after collection and concentration. MS ( $ES^+$ ):  $m/z$  ( $M+Na$ ) = 601.3

This hard white foam (0.16 g, 0.28 mmol) in anhydrous dichloromethane (3 mL) is cooled to 0 °C in an ice-water bath and hydrochloric acid (4 N solution in dioxane, 1 mL, 4 mmol) is added. Stirring at 0 °C is continued for 5 minutes and then the cooling bath is removed. After 1 hour, TLC revealed a preponderance of starting material. An additional 2 mL of 4N hydrochloric acid is added. After 2 hours, the reaction mixture is still composed of mostly starting material. The reaction mixture is left in a -10 °C freezer for 72 hours, following which the TLC showed no change. An additional 2 mL of 4 N hydrochloric acid is added and the reaction mixture is allowed to stir for 8 hours at room temperature. LC/MS analysis showed the conversion to product to be nearly complete. Stirring is continued overnight and after which the ratio is unchanged. Solvent and




excess acid are removed under reduced pressure. The resulting light beige foam is triturated with ether, but a gum resulted. Under vacuum, the gum afforded (0.17 g, > 100 %) a light beige foam, which is carried on to the next step without further purification.

To a solution of N, $\beta$ , $\beta$ -trimethyl-L-phenylalanine (0.17 g, 0.56 mmol, Compound 5 15) and the hydrochloride from the previous step (0.28 mmol max) in anhydrous dimethylformamide (4 mL) under a nitrogen atmosphere, hydroxybenzotriazole (0.076 g, 0.56 mmol, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.11 g, 0.56 mmol) and N-methylmorpholine (0.62  $\mu$ L, 0.56 mmol) are added. After 24 hours the mixture is diluted with water, and the aqueous layer is extracted with diethyl ether (3 10 times). The combined organic extracts are washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride, dried over sodium sulfate, filtered and concentrated *in vacuo*. The crude residue is isolated as cloudy beige semi-solid (0.36 g). MS ( $ES^+$ ):  $m/z$  ( $M+Na$ ) = 790.5

A solution of this material (0.36 g, 0.28 mmol maximum) in dichloromethane (5 15 mL) is cooled to 0 °C in an ice-water bath. Trifluoroacetic acid (1.5 mL) is added and the mixture is stirred for 10 minutes at 0 °C and then the cooling bath is removed. After an additional 2 hours of stirring, thin-layer chromatography (TLC) showed complete deprotection. Solvent and excess acid are removed under reduced pressure to yield 0.52 g of a reddish oil. The oil is taken up in tetrahydrofuran (2 mL), methanol (2 mL), and 20 water (1 mL). Lithium hydroxide monohydrate (35 mg, 0.84 mmol) is added and the reaction mixture is stirred at 45 °C for 2 hours. LC/MS analysis showed only slight hydrolysis of the ester. An additional 20 mg of lithium hydroxide monohydrate is added and the reaction mixture is heated at 55 °C overnight. The solvent is evaporated under reduced pressure. The residue is purified by semi-preparative reverse-phase HPLC 25 (employing a gradient elution of 5 % acetonitrile/95 % water/0.1 % trifluoroacetic acid to 100 % acetonitrile over 1 hour) to give N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N, $\beta$ , $\beta$ -trimethyl-L-phenylalaninamide trifluoroacetic acid as a hard, white foam (0.12 g, 50 % over 4 steps). TOF MS ( $ES^+$ ):  $m/z$  ( $M+H$ ) = 640.4

30

**TABLE 1: Photoaffinity Probes for HTI-286**

Compound	Structure	$K_D$ ( $\mu M$ ) <sup>†</sup>	$IC_{50}$ (nM) †	tubulin polymerization (% inhibition) †
HTI-286		0.4 – 0.8	$0.96 \pm 0.5$ (n = 79)	$87.5 \pm 12.2$ (n = 49)
Probe 1		0.2 – 0.8	$1.8 \pm 0.1$ (n = 2)	88
Probe 2		1.1 - 6	$22.4 \pm 0.7$ (n = 2)	69

<sup>†</sup> Data are from experiments described in Section 7.3, below.

#### 7.2.4 Radiolabeling of Probe 1 and 2

Probes 1 and 2 were tritium labeled by direct tritium exchange labeling of the corresponding unlabeled compounds using Crabtree's catalyst, (1,5-cyclooctadiene)(pyridine)(tricyclohexylphosphine)iridium(I) hexafluorophosphate (Crabtree *et al.*, *J. Am. Chem. Soc.* 1982, 104:6994-7001; Crabtree, *Acc. Chem. Res.* 1990, 23:95-101). This catalyst is commonly used for the introduction of tritium labels into aromatic positions, especially those positions *ortho* to benzophenone carbonyl groups (Heys *et al.*, *J. Chem. Soc. Chem. Commun.* 1992, 9:680-681; Hesk *et al.*, *J. Label Compd. Radiopharm.* 1995, 36:497-502; Shu *et al.*, *J. Organometallic Chem.* 1996, 524:87-93; Chen *et al.*, *J. Label Compd. Radiopharm.* 1997, 39:291-298; Shu *et al.*, *J. Label Compd. Radiopharm.* 1999, 42:797-807). A solution of unlabeled Probe 1 TFA salt (1 mg, 1.6  $\mu$ mol) and Crabtree's catalyst (5 mg, 6.4  $\mu$ mole, 4.0 equiv; purchased from Aldrich (St. Louis, MO)), in methylene chloride (1.00 mL) was exposed to 3.90 Curies of pure tritium gas (0.20 atmosphere) overnight followed by work-up, removal of all volatile tritium, and HPLC purification on a Phenomenex Prodigy 5 $\mu$ m ODS(3) semi-preparative HPLC column (250mmL x 10mm ID) with an aqueous acetonitrile (0.02% TFA)

gradient. A total of 89.9 mCi of tritiated Probe 1 was recovered with radiochemical purity > 98%. The specific activity was determined to be 70.6 Ci/mmol by LC/MS, with an average of 2.45 tritium atoms per molecule. Proton (600 MHz) and tritium (640 MHz) NMR (in  $d_6$ -DMSO solvent) established that the labels were in aromatic positions as expected. Similarly, a solution of unlabeled Probe 2 bis TFA salt (1.4 mg, 1.6  $\mu$ mole) and Crabtree's catalyst (6.5 mg, 8.0  $\mu$ mole, 5.0 equiv) in methylene chloride (1.00 mL) was exposed to 3.79 Curies of pure tritium gas (0.21 atmosphere) overnight followed by work-up, removal of all volatile tritium, and HPLC purification as described above. A total of 93.5 mCi of tritiated Probe 2 was recovered with a radiochemical purity > 98%. The specific activity of tritiated Probe 2 was 82.1 Ci/mmol as determined from LC/MS, with an average of 2.80 tritium atoms per molecule. Similar proton and tritium NMR analysis (also in  $d_6$ -DMSO solvent) confirmed the presence of the tritium atoms in aromatic positions.

### 7.3. Binding and Activity Studies of HTI Analog Probes

This example describes binding and activity studies of HTI analogs that are referred to here as Probes 1 and 2, and are described in the Examples at sections 7.2.1 and 7.2.2, *supra*. The results of these studies, including binding affinity, tubulin polymerization, and cell proliferation assays, demonstrate that the HTI analogs of this invention interact with tubulin and have other properties similar to HTI-286. The results additionally demonstrate that these analogs inhibit tubulin polymerization with a potency that is at least comparable to the potency reported for HTI-286 and for other anti-microtubule agents (Hamel, *Med. Res. Rev.* 1996, 16:207-231).

#### 7.3.1 Reversible binding affinity

Using the fluorescence method described *supra*, the apparent binding constants ( $K_D$ ) for Probes 1 and 2 were determined to be 0.5 and 3.6  $\mu$ M, respectively. The  $K_D$  obtained for Probe 1 is approximately equivalent to the  $K_D$  value of 0.6  $\mu$ M obtained for HTI-286. The  $K_D$  obtained for Probe 2 is approximately 6-fold higher than the  $K_D$  value obtained for HTI-286. These results suggest that Probe 1 and Probe 2 have binding affinities for tubulin that are at least comparable to the affinity of their parent compound HTI-286.

Applying the radiolabeled probe technique described *supra*, an apparent  $K_D$  value for Probe 1 of 0.35  $\mu$ M was determined. This result, which is in close agreement with the

value determined in the fluorescence binding assay (*supra*), further verifies that Probe 1 and HTI-286 have similar binding affinities.

### 7.3.2 Tubulin polymerization in a cell-free system

5       As described *supra*, tubulin polymerization in a cell-free system was performed in the presence or absence of test compound to determine the extent of tubulin polymerization inhibition by test compounds. Consistent with reversible binding affinity results, the compounds were good inhibitors of microtubule-mediated phenomena according to the above assay. In particular, 0.3  $\mu$ M HTI-286, Probe 1, and Probe 2  
10       inhibited tubulin polymerization in a cell-free system by 88%, 88%, and 69%, respectively.

### 7.3.3 Cell proliferation in tissue culture

      According to methods *supra*, cell proliferation in tissue culture was determined in  
15       the presence or absence of test compound to determine the extent of cell proliferation inhibition by test compounds. Probes 1 and 2 inhibit the proliferation of KB cells in tissue culture with potency values ( $IC_{50}$  = 1.8 nM and 22 nM, respectively) that are comparable to the potency of HTI-286 ( $IC_{50}$  = 0.96 nM)

## 20       7.4. Photolabeling of Tubulin using Hemiasterlin Analog Probe 1

### 7.4.1 Probe 1 photolabels the $\alpha$ -tubulin subunit

      The binding of Probe 1 to tubulin was evaluated by incubating tubulin derived from bovine brain (2.5  $\mu$ M) or HeLa cells (10  $\mu$ M) with 2.5  $\mu$ M (3.6  $\mu$ Ci) of [ $^3$ H]-Probe 1  
25       for 30 minutes at room temperature followed by irradiation at 360 nm for two hours at 4°C. Two hours of irradiation was chosen since initial experiments indicated that photoincorporation reached maximum levels at this time.

      As noted above, irradiation of the benzophenone derivative probes at this wavelength effectively activates the photoprobe with minimal destruction of the protein.  
30       At the same time, and in contrast to other photolabeling compounds (*e.g.* with azido-containing photoprobes), irradiation of the benzophenone derivative probes used here leads to an excited state that reverts back to the ground state in the absence of an abstractable proton. This property, combined with the short lifetime of the excited

state(s) of these probes, is understood to preclude the possibility of "false labeling" (*i.e.* of cross-linking with a region of the protein that is not a true labeling site).

5  $\alpha$  and  $\beta$ -tubulin were resolved by SDS-PAGE and analyzed by Coomassie blue staining or fluorography of the gel as described above. According to **Figure 1A**, the Coomassie blue staining reveals an  $\alpha$ - and  $\beta$ -tubulin band for both the bovine and Hela cell tubulin used in this example. In contrast, the autoradiograph revealing radioactive, cross-linked probe yields a band for only the  $\alpha$ -tubulin derived from human and bovine sources. Hence, Probe 1 binds exclusively to  $\alpha$ -tubulin. Similar results are observed when the irradiation time is varied between 5 minutes and 4 hours. Any labeling  
10 observed to co-migrate with  $\beta$ -tubulin is detected only after the species labeling  $\alpha$ -tubulin is grossly overexposed and can not be definitively distinguished from  $\alpha$ -tubulin. The identification of the higher of the two molecular weight species within the gels can be confirmed to be  $\alpha$ -tubulin using monoclonal antibodies specific for  $\alpha$  and  $\beta$ -tubulin, as well as by mass spectrometric analysis of the  $\alpha$  and  $\beta$ -tubulin bands.

15 Probe 1 also specifically labels tubulin when co-incubated with cytosolic preparations from cell lysates that contain numerous other proteins (**Figure 1B**). As shown in lane 1 of the fluorograph in **Figure 1B**, non-radiolabeled Probe 1 can inhibit the binding of radiolabeled Probe 1 in these lysates.

In addition, Probe 1 specifically labels tubulin when incubated in the presence of  
20 whole KB-3-1 cells. The fluorograph pertaining to this experiment reveals two bands for the sample incubated only with radiolabeled Probe 1, a 50 kDa band that co-migrates with purified tubulin and a 65 kDa band. In a sample pre-incubated with non-radiolabeled HTI-286 before exposure to radiolabeled Probe 1, the fluorograph shows only the band at 65 kDa. Thus, competition with non-radiolabeled HTI-286 eliminates the specific  
25 labeling of the 50 kDa tubulin band. Bands in the fluorographic image at 65 kDa show that radiolabeled Probe 1 non-specifically labels cells, since this band remains in the fluorograph when the sample is pre-incubated with non-radiolabeled HTI-286.

30 7.4.2 HTI-286 inhibits the photolabeling of tubulin by Probe 1 further confirming that Probe 1 acts in a similar manner compared to its parent compound

The specificity of photolabeling by Probe 1 was demonstrated by pre-incubating bovine brain tubulin (2.5  $\mu$ M) with 1 mM HTI-286 for 15 minutes at 4°C, followed by addition of increasing concentrations of radiolabeled Probe 1 (0.025-2.5  $\mu$ M) as described



above. After incubation for 30 minutes at 37°C, samples were irradiated with 360 nm UV light for 30 minutes and separated by SDS-PAGE. Gels were exposed to film using fluorographic techniques. Under these conditions, 1 mM HTI-286 completely inhibits the incorporation of 0.25  $\mu$ M radiolabeled Probe 1 into  $\alpha$ -tubulin (Figure 2, compare Lanes 3 and 5) and inhibits the incorporation of 2.5  $\mu$ M radiolabeled Probe 1 into  $\alpha$ -tubulin by more than 50% (Figure 2, compare Lanes 4 and 6).  $\alpha$ -tubulin was the only species labeled and no detection of  $\beta$ -tubulin labeling was found under any condition confirming that Probe 1 binds exclusively to  $\alpha$ -tubulin.

Photolabeling specificity for Probe 1 was also confirmed by labeling tubulin in the presence of increasing concentrations of radiolabeled Probe 1 in the presence or absence of 100  $\mu$ M non-radiolabeled probe 1. When a ratio of tubulin : radiolabeled probe 1 of 2.5:0.25 (10:1) was used, tubulin labeling was inhibited 93% by non-radiolabeled probe 1. When the ratio was changed to 2.5:2.5 (1:1), conditions used mostly for peptide mapping studies, tubulin labeling was inhibited 82%.

#### 7.4.3 Temperature Effects of Probe 1 Photolabeling of Tubulin

Because the polymerization state of tubulin can be affected by temperature (for example, see Johnson and Borisy, *J. Mol. Biol.* 1979, 133:199-216), experiments were also performed to determine whether the binding of photolabeling probes, such as Probe 1, is also affected by temperature. In particular, photolabeling experiments were performed as described in section 7.1.5 above, at both room temperature and at 37°C. These experiments were conducted by pre-incubating tubulin in either the presence or absence of various competitors (including non-radiolabeled Probe 1, dolastatin-10, HTI-286, vinblastine, paclitaxel, or colchicine) before incubation with radiolabeled Probe 1. Results from these experiments are summarized in Table 2, *infra*, at the end of this Section.

Initial incubations were done at RT using 2.5  $\mu$ M tubulin and 0.25  $\mu$ M radiolabeled Probe 1. Under these conditions, a 400-fold molar excess (100  $\mu$ M) of either non-radiolabeled Probe 1 or dolastatin-10 reduces labeling by approximately 87% whereas either HTI-286 or vinblastine reduces labeling by approximately 72%. In contrast, pre-incubation with either paclitaxel or colchicine has almost no effect on the observed photolabeling of tubulin (Table 2).

When incubations are performed at 37°C, non-radiolabeled Probe 1 and dolastatin-10 are still the best inhibitors and both reduce labeling by about 75%. HTI-286 and vinblastine both reduce labeling by about 50%. However, paclitaxel and colchicine actually enhance labeling by 73 and 119%, respectively, at this temperature -- in contrast with the effect of those compounds at room temperature. The unusual effect of colchicine on Probe 1 binding to tubulin was further investigated by pre-incubating tubulin with 0, 200, 400, 1000, and 4000-fold molar excess of colchicine prior to labeling tubulin with Probe 1 at 37°C. Under these conditions, colchicine maximally enhances binding of the probe when used at a 400-fold molar excess (100 µM), has less enhancing effects at a 1000-fold molar excess (250 µM), and has no effect when used at a 4000-fold excess (1 mM).

**TABLE 2: Competition of Probe 1 Binding to Tubulin**  
**Inhibition (%)**

Competitor	Room Temp.	37 °C
Control (no competitor)	100	100
Probe 1	12 ± 7.8	24 <sup>†</sup>
Dolastatin-10	13 ± 3	26 <sup>†</sup>
HTI-286	27 ± 4	47 <sup>†</sup>
Vinblastine	28 ± 1	52 ± 0.5
Paclitaxel	83 ± 41	173 ± 7
Colchicine	118 ± 17	219 ± 14

<sup>†</sup> Results from a single experiment.

#### 7.4.4 Inhibition of Probe 1 photolabeling by various compounds

The potency of the inhibitory effect of dolastatin-10, HTI-286, and vinblastine on photolabeling was further investigated by determining IC<sub>50</sub> values for these candidate inhibitors (*i.e.* the concentration of these compounds that inhibit Probe 1 photolabeling of tubulin by 50%). The same reaction conditions, using incubation at room temperature, was done as described in Section 7.4.3, above. Under these conditions, Probe 1 and dolastatin-10 are equally potent inhibitors of tubulin photolabeling (IC<sub>50</sub> = 5 µM). HTI-286 is slightly less potent (IC<sub>50</sub> = 7 µM), while vinblastine is the least inhibitory of these compounds (IC<sub>50</sub> = 22 µM).

#### 7.4.5 Effect of GTP on Probe 1 photolabeling of tubulin

It has been previously reported that hemiasterlin and dolastatin-10 interact with tubulin to inhibit GTP exchange, but do not inhibit the ability of GTP to bind to the protein (see, for example Bai *et al.*, *Biochemistry* 1999, 38:14302-14310; Bai *et al.*, *J. Biol. Chem.* 1990, 265:17141-17149). Therefore experiments were also performed to determine whether GTP inhibits binding of Probe 1 to tubulin. In particular, tubulin samples were pre-incubated with a range of different GTP concentrations (between 0 and 100  $\mu$ M) before photolabeling with Probe 1 as described above. However, GTP did not significantly inhibit incorporation of this probe, further indicating that Probe 1 of this invention interacts with tubulin via the same mechanism as its parent hemiasterlin derivative.

#### 7.5. Identification of the Hemiasterlin Binding Site in Tubulin using Probe 1

The experiments described, *supra*, demonstrate that the HTI analogs of this invention (*e.g.* Probes 1 and 2) bind tubulin with properties that are similar to hemiasterlin. Accordingly, this section describes experiments that identify the binding site of hemiasterlin and/or other hemiasterlin analogs (for example, HTI-286) to tubulin. In particular, experiments are described using radiolabeled and/or photo-probe of the invention, Probe 1, to identify the tubulin binding site. The cross-linking efficiency for Probe 1 was approximately 1.5% under optimal conditions and found to be sufficient for peptide mapping studies.

##### 7.5.1 Formic Acid digestion after Probe 1 photoaffinity labeling

To ascertain an initial labeling region for Probe 1 in  $\alpha$ -tubulin, formic acid digestions were performed. Formic acid is known to preferentially cleave Asp-Pro bonds (Sonderegger *et al.*, *Anal. Biochem.* 1983, 122:298-301).  $\alpha$ -tubulin is extraordinarily conserved across vertebrate species, including those derived from porcine, rat, murine, and human origin, and contains only one such linkage at Asp<sup>306</sup>-Pro<sup>307</sup> (Figure 3A). Therefore, complete formic acid digestion of  $\alpha$ -tubulin derived from bovine tubulin would be expected to produced two distinct peptide fragments consisting of amino acids 1-306 (~34.5 kDa) and 307-451 (~16 kDa) as obtained from the digestion of other vertebrate species (*e.g.* the species whose tubulin amino acid sequences are depicted in Figures 12A-12B). These fragments have been previously identified by immunological

methods as the N- and C-terminus portions of tubulin, respectively (Chau *et al.*, *Biochemistry* 1998, 37:17692-17703). In contrast, formic acid digestion of  $\beta$ -tubulin is known to produce three fragments consisting of amino acids 1-31 (~3.5 kDa), 32-304 (~31 kDa) and 305-445 (~16 kDa) (Hall *et al. Mol. Cell Biol.* 1983, 3:854-862). Upon  
5 formic acid in-gel digestion of photolabeled tubulin as described in Section 7.1.6, three main protein bands were observed (**Figure 3B**): the upper approximately 34.5 kDa band corresponding to a fragment consistent with  $\alpha$ -tubulin digestion, the middle band at approximately 30 kDa originating from the  $\beta$ -tubulin digestion, and a strongly-staining Coomassie band at approximately 16 kDa corresponding to two 16 kDa fragments  
10 originating from the  $\alpha$  and  $\beta$ -tubulin digestions. Radiolabel derived from Probe 1 was found solely in the 16 kDa peptide fragment (**Figure 3C**). This results is in contrast to a previously described paclitaxel analog that labeled the 31 kDa peptide fragment derived from  $\beta$ -tubulin (Rao *et al., J. Bio. Chem.* 1995, 270:20235-20238). Mass spectrometric methods confirmed that the labeled 16-kDa formic acid fragment was from  $\alpha$ -tubulin.

15

#### 7.5.2 Trypsin digestion of native tubulin after Probe 1 photoaffinity labeling

To define more narrowly the binding region of hemiasterlin and its analogs, trypsin digestion of native tubulin was performed after Probe 1 photoaffinity labeling as described in Section 7.1.7. Under these conditions and due to the high sequence  
20 conservation across vertebrate species, it can be predicted that trypsin preferentially cleaves  $\alpha$ -tubulin after amino acid residue 339, producing two fragments that migrate at approximately 38 and 17 kDa (**Figure 4A**) (Sacket and Wolff, *J. Biol. Chem.* 1986, 261:9070-9076; Serrano *et al., J. Biol. Chem.* 1984, 259:6607-6611; Mandelkow *et al., J. Mol. Biol.* 1985, 185:311-327). The latter fragment has a predicted molecular weight of  
25 14 kDa, but migrates aberrantly in gels (Chau *et al., Biochemistry* 1998, 37:17692-17730). In the experiments, photolabeled tubulin was digested for 5, 10, and 30 minutes with trypsin and the resultant fragments were resolved in gels under conditions that allow  $\alpha$ - and  $\beta$ -tubulin to co-migrate. Both expected species were observed by analyzing the digest using Coomassie blue staining (**Figure 4B**). The origin of the 34 - and 21- kDa  
30 Coomassie blue bands believed to be derived from  $\beta$ -tubulin (Sacket and Wolff, *J. Biol. Chem.* 1986, 261:9070-9076). In contrast to the Coomassie stained bands, the major radiolabeled species are detected at approximately 51 kDa and approximately 38 kDa and co-migrate with the Coomassie-stained species (**Figure 4C**). There is a precursor-product

relationship between the undigested tubulin (51 kDa) and the 38 kDa-species, such that the ratio between the 51 and 38 kDa species decreases as the incubation time with trypsin increases. Both the gel image showing radiolabeled fragments and the radioactivity profile for the gel demonstrate this relationship (**Figure 4C and 4D**, respectively). The origin of a 19 kDa radiolabeled species, which was detected in the autoradiogram of the gel (**Figure 4C**) and slices of gel (**Figure 4D**), is unknown. However, it was present in the undigested sample and did not increase in intensity as the time of trypsin digestion was increased from 5-30 minutes. This may be an inherent fragment in the tubulin preparation. A minor radiolabeled species that migrated at approximately 34 kDa was observed after 10 and 30 minute digestion with trypsin. It is unlikely that this corresponds to the 34 kDa derived from  $\beta$ -tubulin, since the 34 kDa Coomassie-stained band appeared within 5 minutes after trypsin digest, but it was not radiolabeled. Rather, it is more likely that the 34 kDa radiolabeled fragment is a partial digest of the 38 kDa fragment derived from  $\alpha$ -tubulin. Consistent with this hypothesis, the intensity of the 34 kDa radiolabeled band increased as the 38 kDa band intensity decreased. Taken together with the formic acid digestion results, these data demonstrate that Probe 1 labels between residues 307-339 of intact  $\alpha$ -tubulin.

#### 7.5.3 CNBr, trypsin, and Lys C cleavage with Mass Spectrometric Confirmation after Photoaffinity Labeling of Tubulin by Probe 1

To resolve further the Probe 1 photoaffinity labeling domain, digestion of Probe 1-labeled tubulin was done with CNBr. CNBr cleaves peptide bonds on the carboxy terminus side of methionine residues and, in these studies, was used to hydrolyze peptide bonds in polyacrylamide gels (Loeb *et al.*, *Anal. Biochem.* 1989, 176:365-367). Based on the sequence conservation of  $\alpha$ -tubulin across a variety of species, it was predicted that  $\alpha$ -tubulin would be digested into 11 fragments by CNBr (**Figure 5C and 5D**). One of these fragments, which is conserved across rat, mouse, and pig  $\alpha$ -tubulin, has a predicted molecular weight of 7 kDa that spanned residues 314 to 376. To determine if the labeling site resided in the 7 kDa CNBr fragment as predicted from the labeling experiments above,  $\alpha$  and  $\beta$ -tubulin were separated in gels and digested with CNBr according to the methods described in Section 7.1.8. Each digestion was separated on a gel and observed either by using fluorography or after silver staining of the gel (**Figures 5A and 5B**, respectively). The numerous silver-stained fragments observed in **Figure 5B** indicate that

the CNBr digestion of  $\alpha$  and  $\beta$ -tubulin occurred. Upon fluorography of the gel, a major and minor radiolabeled species were observed at 7 and 8 kDa, respectively (**Figure 5A**). Mass spectrometric results confirm that the 7 kDa fragment was correctly identified compared to the predicted results. The origin of the 8 kDa species is unknown, but likely to be a partial digest composed of the 1291 and 7166 dalton CNBr fragments (**Figure 5D**). These cleavage studies further defined the binding site of Probe 1 to reside between residues 314 and 339.

#### 10                    7.5.4   Effects of the Labeling of $\alpha$ -Tubulin by Probe 1 on Subtilisin Digestion.

To ensure that a major labeling site was not overlooked, studies were done with subtilisin digestion before or after Probe 1 photolabeling of tubulin as described in Section 7.1.10. Knowing that  $\alpha$ -tubulin is conserved across many species, it can be inferred that susubtilisin cleaves bovine  $\alpha$ -tubulin between Asp-438 and Ser-439 and  $\beta$ -tubulin between Gln-433 and Gly-434 (Redeker *et al.*, *FEBS Lett.* 1992, 313:185-192) (**Figure 6A**). The cleaved protein is referred to as tubulin-S (Serrano *et al.*, *Proc. Natl. Acad. Sci. USA* 1984, 81:5989-5993). Consistent with these predictions, it is observed that subtilisin digestion increases the mobility of  $\alpha$ - and  $\beta$ -tubulin approximately 2 kDa as detected by Commassie staining of the material resolved in gels (**Figure 6B**, compare Lanes 1 and 2, upper panel). If enzyme digestion is done before labeling, a radiolabeled band that co-migrates with the position of  $\alpha$ -tubulin-S is detected (**Figure 6B**, Lane 2). Since the intensity of radiolabeled  $\alpha$ -tubulin-S does not decrease remarkably compared with non-digested material, especially when considering the relative amounts of Commassie-staining material that are detected, this suggests that a major photolabeling domain is unlikely to reside in the C-terminus of  $\alpha$ -tubulin. This conclusion is consistent with the findings stated above.

To eliminate further the possibility of a C-terminus binding domain for Probe 1, the C-terminus of  $\alpha$ -tubulin was removed by subtilisin digestion after photolabeling (**Figure 6B**, Lane 3). Contrary to the expected result,  $\alpha$ -tubulin-S is not labeled with probe 1, despite the fact that digestion did occur according to Coomassie staining. Some radiolabeled tubulin remains at the position of  $\alpha$ -tubulin. Since vinblastine (but not maytansine) has been shown to inhibit the digestion of  $\alpha$ -tubulin but not  $\beta$ -tubulin by subtilisin, and this effect may be mediated by alteration of the state of tubulin rather than

blockade of the cleavage site by vinblastine (Rai and Wolff, *Proc. Natl. Acad. Sci. USA* 1998, 95:4253-4257), it is hypothesized that hemiasterlins perform in a similar fashion.

Upon closer inspection, it was found that 0.25  $\mu$ M radiolabeled [ $^3$ H]-Probe 1 partially inhibits the digestion of  $\alpha$ -tubulin (**Figure 6C**, Lane 2, Coomassie stained gel) and 100 or 250  $\mu$ M non-radiolabeled Probe 1 completely blocks the ability of  $\alpha$ -tubulin to be digested with subtilisin, while  $\beta$ -tubulin is completely digested (**Figure 6C**, Lanes 3-6). These results suggest that the C-terminus of  $\alpha$ -tubulin and the hemiasterlin photolabeling site interact such that the C-terminus cannot be digested with subtilisin if tubulin is pre-incubated with Probe 1. However as observed above, the C-terminus of  $\alpha$ -tubulin does not contain a major labeling site.

## 7.6. Photolabeling of Tubulin using Hemiasterlin Analog Probe 2

### 7.6.1 Probe 2 photolabels the $\alpha$ -tubulin subunit

Probe 2 was observed to photolabel tubulin using the methods described in Section 7.1.5. Similar to Probe 1, Probe 2 exclusively labels  $\alpha$ -tubulin in purified tubulin preparations from bovine brain and HeLa cells (**Figure 7**). In contrast to the Coomassie blue stained gel (**Figure 7A**) that displays both  $\alpha$  and  $\beta$ -tubulin, the autoradiograph shows that only  $\alpha$ -tubulin contains radiolabeled photoprobe (**Figure 7B**).

### 7.6.2 Competition Studies using Various Anti-cancer Drugs as Inhibitors of Probe 2 Photolabeling

Competition binding experiments were performed as described in Section 7.1.5. Briefly, 100  $\mu$ M competitor drug was pre-incubated at room temperature for 15 minutes with 2.5  $\mu$ M tubulin before incubating with 0.25  $\mu$ M [ $^3$ H]-Probe 2 at room temperature for 30 minutes. Samples were then irradiated for 2 hours at 4°C and evaluated by SDS-PAGE and fluorography. Non-radioactive Probe 2, dolastatin-10, HTI-286 and vinblastine can inhibit Probe 2 labeling of tubulin by 12%, 27%, 45%, and 64%, respectively. As seen in the case of Probe 1, paclitaxel and colchicine enhanced photolabeling to 113% and 107% of the control value, respectively. However, this effect was lost at higher concentrations of paclitaxel and colchicine. The ability for certain competitors to reduce or increase Probe 2 labeling demonstrates that Probe 2 specifically labels tubulin.

### 7.7. Identification of the Hemiasterlin Binding Site in Tubulin using Probe 2

The experiments described, *supra*, demonstrate that the HTI analogs of this invention (*e.g.* Probes 1 and 2) bind tubulin with properties that are similar to hemiasterlin. Accordingly, this section describes experiments that identify the binding site of hemiasterlin and/or other hemiasterlin analogs (for example, HTI-286) to tubulin. In particular, experiments are described using the radiolabeled photo-probe of the invention, Probe 2, to identify the tubulin binding site. These studies can be paired with Probe 1 mapping studies described in Section 7.5 to define further the binding site of hemiasterlin and its analogs.

#### 7.7.1 Trypsin, Formic Acid, and Lys C digestion after Probe 2 photoaffinity labeling

To determine the Probe 2 labeling site on  $\alpha$ -tubulin, trypsin, formic acid, and Lys C digestions were performed on photolabeled bovine brain tubulin as described in Sections 7.1.7 and 7.1.11, respectively. As described in Section 7.5.2 and **Figure 4A**, trypsin cleaves  $\alpha$ - and  $\beta$ -tubulin partially under native conditions. The first fragments obtained are 38 kDa and 17 kDa for  $\alpha$ -tubulin and 34 kDa and 21 kDa for  $\beta$ -tubulin. Probe 2 was found to be bound to the 38 kDa N-terminal fragment from  $\alpha$ -tubulin spanning residue 1-338 (**Figure 8A and 8B**).

Formic acid cleavage was carried out in the presence of SDS as a denaturant such that all formic acid sites for  $\alpha$ - and  $\beta$ -tubulin are accessible and digestion is complete. As described in Section 7.5.1, formic acid digestion results in the cleavage of  $\alpha$ -tubulin into two fragments 34.5 kDa (residues 1-306) and 16 kDa (307-451) (**Figure 3A**). In contrast, formic acid digestion of  $\beta$ -tubulin is known to produce three fragments consisting of amino acids 1-31 (3.5 kDa), 32-304 (31 kDa) and 305-445 (16 kDa). The Probe 2 radiolabel was found to label the 34.5 kDa fragment that corresponds to  $\alpha$ -tubulin residues 1-306 (**Figure 8A and 8B**).

In a separate digestion experiment, Lys C was used to digest both  $\alpha$ - and  $\beta$ -tubulin into several fragments. As predicted from the sequence homology with other vertebrate species, the largest bovine brain  $\alpha$ -tubulin fragment was 12.8 kDa (**Figure 8E**). Probe 2 was found to label this 12.8 kDa fragment corresponding to residues 167-280 (**Figure 8C**



and 8D). Combining the results for the trypsin, formic acid, and Lys C digestions, Probe 2 photolabels  $\alpha$ -tubulin between residues 167-280.

#### 7.7.2 Subtilisin digestion before Probe 2 photoaffinity labeling

5 To define further the photolabeling and binding site of Probe 2, subtilisin digestions were performed before Probe 2 photoaffinity labeling as discussed in Section 7.1.12. Subtilisin cleaves a 2-4 kDa C-terminal fragment from both  $\alpha$ - and  $\beta$ -tubulin as described in Section 7.5.4 and **Figure 6A**. After cleavage of native tubulin with subtilisin and [ $^3\text{H}$ ]-Probe 2 photolabeling, most of the label was present in the 48 kDa tubulin  
10 fragment (**Figure 8A and 8B**). Thus, the 48 kDa fragment contains the binding region for probe 2. This result further verifies that the binding and labeling site for Probe 2 defined by the trypsin, formic acid, and Lys C digestions is valid.

#### 7.7.3 CNBr cleavage after Photoaffinity labeling of Tubulin by Probe 2

15 To determine more narrowly the binding site region for Probe 2, CNBr digestions according to the methods described in Section 7.1.8 were performed. As discussed in Section 7.5.3 and **Figure 5**, CNBr digestion of  $\alpha$ - and  $\beta$ -tubulin also gives rise to several fragments of approximately equal molecular weight. To minimize the number of probable fragments and their identity, photolabeled tubulin was separated into  $\alpha$ - and  $\beta$ -  
20 tubulin subunits by SDS-PAGE, and the bands corresponding to the two proteins were digested separately with CNBr. Digested  $\alpha$  and  $\beta$ -tubulin were run on gels separately and either silver stained (**Figure 9A**) or subjected to fluorographic methods (**Figure 9B**) as described in Section 7.1.8. The radiolabeled band with the highest intensity corresponds to a 12 kDa fragment (residues 204-302). Other bands that contain radiolabel are located  
25 at 13kD and 20kDa (**Figure 9B**) and possibly correspond to partial digests with CNBr at residues 204-313 and 204-377 respectively. Taken together with other digests described in this section, these results suggest a binding site for Probe 2 on  $\alpha$ -tubulin within residues 204-280.

#### 30 7.8. Subtilisin Cleavage Studies for Tubulin in the presence of various anti-microtubule agents

Vinblastine protects against subtilisin cleavage of  $\alpha$ -tubulin as described in Section 7.5.4 (Rai & Wolff, *Proc. Natl. Acad. Sci. U.S.A.* 1998, 95:4253-4257). The

hemiasterlin analog, HTI-286, was compared with other microtubule binding agents to determine if HTI-286 also protects  $\alpha$ -tubulin against subtilisin cleavage and if this protection occurs through the same proposed mechanism as other binding agents.

Methods for this experiment are described in Section 7.1.13. Partial cleavage of  $\alpha$ - and  $\beta$ -tubulin by subtilisin gives rise to a 2-4 kDa shift of the two subunits as seen in SDS-PAGE gels (**Figure 10**, Control). If the amount of subtilisin and time of cleavage is controlled, the  $\beta$ -tubulin is cleaved more rapidly than  $\alpha$ -tubulin. When tubulin was incubated with HTI-286, dolastatin-10, vinblastine and paclitaxel, subtilisin was unable to cleave  $\alpha$ -tubulin even after 60 minutes (**Figure 10**) as seen by the lack of shifting of the  $\alpha$ -tubulin as compared to the Control lane. Colchicine was the only drug that did not protect  $\alpha$ -tubulin from cleavage by subtilisin.

The protection of  $\alpha$ -tubulin by vinblastine has been attributed to the ability of drugs to induce the formation of microtubules (in case of paclitaxel) or tubulin polymers/aggregates (in the case for vinblastine and dolastatin-10). See, Rai & Wolff, *Proc. Natl. Acad. Sci. U.S.A.* 1998, 95:4253-4257. These aggregates can be recovered by centrifugation at high speeds as described in Section 7.1.14. When drug treated tubulin was subjected to centrifugation at 10,000 g and 100,000 g a significant amount of protein was recovered in the pellet fractions of tubulin treated with dolastatin-10, vinblastine and paclitaxel, but no significant protein was recovered in HTI-286 and colchicine fractions. Absence of pellet fractions suggests that HTI-286 does not cause aggregation of tubulin, the aggregates are not large enough, or the aggregates are unstable. Nevertheless, despite the lack of stable aggregates, HTI-286 protects  $\alpha$ -tubulin from subtilisin cleavage. This result suggests that the mechanism of protection of  $\alpha$ -tubulin from subtilisin cleavage by HTI-286 binding is distinct from vinblastine and dolastatin-10.

#### 7.9. Mapping the photoaffinity labeling sites for Probe 1 and Probe 2 to the electron micrographic crystal structure of tubulin (3.5 Å)

Molecular modeling studies were performed using the Weblab Viewer Lite software. The structure of tubulin was taken from Lowe *et al.* (*J. Mol. Biol* 2001, 313:1045-1057). The major photoaffinity binding site for Probe 1 is located within residues 314-338 that, based on the electron micrographic crystal structure (3.5 Å) of zinc-induced tubulin sheets (Lowe *et al.*, *J. Mol. Biol* 2001, 313:1045-1057), corresponds to the S8 (residues 311-321) to H10 (residues 324-336) regions of  $\alpha$ -tubulin (**Figure 11**,

green label). The region is perfectly conserved within porcine, rat and murine species (Stanchi *et al.*, *Biochem. Biophys. Res. Commun.* 2000, 270:1111-1118, See, also, **Figures 12A-12B**) with the exception of human  $\alpha$ -tubulin that contains a single amino acid substitution (Met<sup>317</sup>). This region is approximately 80% divergent from the corresponding region in  $\beta$ -tubulin within these species. The H10 domain is involved in establishing both longitudinal and lateral protofilament contacts of tubulin subunits (Nogales *et al.*, *Cell* 1999, 96:79-88). Further resolution of the exact binding site within the S8-H10 region are required to refine the model, but due to the paucity of labeling we have not been able to resolve the site further.


The Probe 2 photoaffinity labeling site exists between residues 204-280 (**Figure 11**, yellow label). The binding site defined by Probe 1 and 2 is a novel site for interaction of anti-mitotic drugs with tubulin. In contrast, the tubulin polymerizer paclitaxel binds in  $\beta$ -tubulin and the tubulin depolymerizing vinca-peptides bind the "vinca binding domain" on  $\beta$ -tubulin. Colchicine binds  $\alpha$ - and  $\beta$ -tubulin in the intradimer interface.

#### 8. REFERENCES CITED

Numerous references, including patents, patent applications and various publications, are cited and discussed in the description of this invention. The citation and/or discussion of such references is provided merely to clarify the description of the present invention and is not an admission that any such reference is "prior art" to the invention described here. All references cited and/or discussed in this specification (including references, *e.g.*, to biological sequences or structures in the GenBank, PDB or other public databases) are incorporated herein by reference in their entirety and to the same extent as if each reference was individually incorporated by reference.

WHAT IS CLAIMED IS:

1. A compound represented by the formula:



or a pharmaceutically acceptable salt thereof, in which:

- (a)  $R_1$  is a photoreactive moiety or an aryl moiety;
- (b)  $R_2$  is a photoreactive moiety, an alkyl moiety or H; and
- (c) at least one of  $R_1$  and  $R_2$  is a photoreactive group.


2. The compound according to claim 1 in which the photoreactive group of  $R_1$  or  $R_2$  is a benzophenone moiety.

3. The compound according to claim 1 in which the photoreactive group of  $R_1$  or  $R_2$  is an azide moiety.

4. A compound according to claim 1 in which  $R_1$  is a benzophenone moiety, and  $R_2$  is an alkyl moiety or H.


5. A compound according to claim 1 in which  $R_1$  is an aryl moiety and  $R_2$  is a photoreactive group.

6. A compound according to claim 1 and having the formula:



or a pharmaceutically acceptable salt thereof.

7. A compound according to claim 1 and having the formula



or a pharmaceutically acceptable salt thereof.

8. A compound according to claim 1 which is 4-benzoyl-N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-N1-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N1,3-dimethyl-L-valinamide or a pharmaceutically acceptable salt thereof.

9. A compound according to claim 1 which is N,β,β-trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N,β,β-trimethyl-L-phenylalaninamide or a pharmaceutically acceptable salt thereof.

10. A method for identifying a tubulin binding site, which method comprises:

- (a) contacting a compound according to claim 1 to a sample comprising tubulin such that the compound is able to irreversibly bind to the tubulin;
- (b) separating the tubulin into a plurality of tubulin fragments; and
- (c) identifying at least one tubulin fragment with the compound bound thereto.

wherein the identification of a tubulin fragment with the compound bound thereto identifies said fragment as a tubulin binding site.

11. The method according to claim 10 in which the sample is irradiated after contacting the compound to the sample.

12. The method according to claim 10 wherein:

- (a) the compound is detectably labeled, and

- (b) a tubulin fragment with the compound bound thereto is identified by detecting the label with said fragment.
13. The method according to claim 10 in which the tubulin is chemically digested.
14. The method according to claim 13 in which the tubulin is chemically digested with formic acid or CNBr.
15. The method according to claim 10 in which the tubulin is enzymatically digested.
16. The method according to claim 15 in which the tubulin is digested with Lys C, Trypsin or subtilisin.
17. The method according to claim 10 in which the compound is 4-benzoyl-N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-N1-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N1,3-dimethyl-L-valinamide or a pharmaceutically acceptable salt thereof.
18. The method according to claim 10 in which the compound is N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N, $\beta$ , $\beta$ -trimethyl-L-phenylalaninamide or a pharmaceutically acceptable salt thereof.
19. A method for identifying a hemiasterlin competitor, which method comprises:
- (a) contacting a test compound and a probe to a sample comprising tubulin, wherein the probe
    - (i) is a compound according to claim 1, and
    - (ii) is contacted to the sample such that it is able to irreversibly bind tubulin;
  - (b) detecting binding of the probe to the tubulin; and
  - (c) comparing said binding to binding of the probe to tubulin in the absence of the test compound,

in which reduced binding in the presence of the test compound identifies said test compound as a hemiasterlin competitor.

20. The method according to claim 19 in which the sample is irradiated after contacting the probe to the sample.

21. The method according to claim 19 in which the test compound is contacted to the sample before the probe.

22. The method according to claim 19 in which the test compound and the probe are contacted to the sample simultaneously.

23. The method according to claim 19 in which the tubulin is digested after being contacted with the probe.

24. The method according to claim 23 in which the tubulin is chemically digested.

25. The method according to claim 24 in which the tubulin is digested with formic acid or CNBr.


26. The method according to claim 24 in which the tubulin is enzymatically digested.


27. The method according to claim 26 in which the tubulin is digested with Lys C, Trypsin or subtilisin.

28. The method according to claim 10 in which the compound is 4-benzoyl-N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-N1-[(1S,2E)-3-carboxy-1-isopropylbut-2-enyl]-N1,3-dimethyl-L-valinamide or a pharmaceutically acceptable salt thereof.


29. The method according to claim 10 in which the compound is N, $\beta$ , $\beta$ -trimethyl-L-phenylalanyl-4-benzoyl-N-[(1S,2E)-3-carboxy-1-isopropyl-2-butenyl]-N, $\beta$ , $\beta$ -trimethyl-L-phenylalaninamide or a pharmaceutically acceptable salt thereof.




**Figure 1A**

**Figure 1B**

<b>Probe 1 (<math>\mu\text{M}</math>) :</b>	0	0.025	0.25	2.5	0.25	2.5
<b>HTI-286 (1 mM) :</b>	-	-	-	-	+	+

**Figure 2**

**Figure 3A**







Figure 3B


**Figure 3C**




**Figure 4A**

**Figure 4B**



**Figure 4C**

**Figure 4D**




Figure 5A




Figure 5B




Figure 5C

**Predicted  
Fragment** →

Mol Wt	Len		10	20	30	40	50	60
149.2	1	H	:	:	:	:	:	:
4027.5	35	RECISIHVGQAGVQIGNACWELYLEHGIQPDGQM ***:					:	:
12994.6	118	PSDKTIGGGDDSFNTFFSETGAGKHVPRAVFVDLEPTVIDEVRTGTYRQLFHPEQLITCKEDAANN-*>						
5562.3	49	ERLSVDYGRKSKLEFSIYPAPQVSTAVVEPYN SILTHTTLEHSDCAFH *						:
11316.7	99	VDNEAIYDICTRRNLDIERPTYTNLNLRLISQIVSSITASLRFDGALNVDLTFQTNLVPYPRIHFPL-*>						
1291.5	11	VKCDPGHGKYM *	:	:	:	:	:	:
7166.4	64	ACCLLYRGDVVPKDVNAAIATIKTKRTIQFVDWCPTGFKVGINYQPPTVVPGDLAKVQRAVCM ****						
2403.7	21	LSNTTATAEAWARLDHKFDLM		:	:	:	:	:
1814.1	15	YAKRAFVHWVVGEGH	:	:	:	:	:	:
1428.4	12	EEGEFSEAREDM	:	:	:	:	:	:
2861.9	26	AALEKDYEEVGVDSVEGEGEEEGEY	:	:	:	:	:	:

Figure 5D




Figure 6A




Figure 6B






Figure 6C




Figure 7A





Figure 7B



**Figure 8A****Figure 8B**

38 kDa →





← 34.5 kDa

Trypsin  
Subtilisin  
Formic Acid

**Figure 8C****Figure 8D**




**Figure 9A**



**Figure 9B**



**Figure 10**




Figure 11

Htub1/1-451	MRECISVHVGQAGVQIGNACWELFCLEHGIQADGTFDAQASKINDDDSFNTFFSETGNGK
Htub2/1-451	MRECISIHVGQAGVQIGNACWELCYCLEHGIQPDGQMPSDKTIGGGDDSFNTFFSETGAGK
ptub1/1-451	-RECISVHVGQAGVQMGNACWELCYCLEHGIQPDGQMPSDKTIGGGDDSFNTFFCETGAGK
ptub2/1-451	-----GQAGVQIGNACWELCYCLEHGIQPDGQMPSDKTIGGGDDSFNTFFSETGAGK
motub1/1-451	MRECISVHVGQAGVQIGNACWELFCLEHGIQADGTFGTQASKINDDDSFNTFFSETGNGK
hamtub1/1-451	MRECISIHVGQAGVQIGNACWELCYCLEHGIQPDGQMPSDKTIGGGDDSFNTFFSETGAGK
rattub1/1-451	MRECISIHVGQAGVQIGNACWELCYCLEHGIQPDGQMPSDKTIGGGDDSFNTFFSETGAGK
chicktub1/1-451	MRECISVHIGQAGVQIGNACWELFCLEHSIQPDGTFSDPPSS---DDSFATFFRETSMK
frogtub1/1-451	MRECISIHVGQAGVQIGNACWELCYCLEHGIQPDGQMPSDKTIGGGDDSFNTFFSETGAGK
Htub1/1-451	HVPRAVMIDLEPTVVDEVVRAGTYRQLFHPEQLITGKEDAANNYARGHYTVGKESIDLVLVD
Htub2/1-451	HVPRAVFVDLEPTVIDEVRTGTyrQLFHPEQLITGKEDAANNYARGHYTIGKEIDLVLVD
ptub1/1-451	HVPRAVFVDLEPTVIDEIRNGPYRQLFHPEQLITGKEDAANNYARGHYTIGKEIDPVLVD
ptub2/1-451	HVPRAVFVDLEPTVIDEVRTGTyrQLFHPEQLITGKEDAANNYARGHYTIGKEIDLVLVD
motub1/1-451	HVPRAVMVDLEPTVVDEVVRAGTYRQLFHPEQLITGKEDAANNYARGHYTVGKESIDLVLVD
hamtub1/1-451	HVPRAVFVDLEPTVIDEVRTGTyrQLFHPEQLITGKEDAANNYARGHYTIGKEIDLVLVD
rattub1/1-451	HVPRAVFVDLEPTVIDEVRTGTyrQLFHPEQLITGKEDAANNYARGHYTIGKEIDLVLVD
chicktub1/1-451	YVPRAIMVDLEPTVVDEVVRAGTYRHLFHPEQLITGKEDAANNYARGHYTVGKDKVDMVSD
frogtub1/1-451	HVPRAVFVDLEPTVIDEVRTGTyrQLFHPEQLITGKEDAANNYARGHYTIGKEIDLVLVD
Htub1/1-451	RIRKLTACSGLQGFLIFHSFGGGTSGSFTSLLMERLSLDYGKSKLEFAIYPAPQVSTA
Htub2/1-451	RIRKLADQCTRLQGFLVFHSFGGGTSGSFTSLLMERLSVDYGKSKLEFSIYPAPQVSTA
ptub1/1-451	RIRKLSAQCTGLQGFLVFHSFGGGTSGSFTSLLMERLSVDYGKSKLEFSIYPAPQVSTA
ptub2/1-451	RIRKLADQCTGLQGFLVFHSFGGGTSGSFTSLLMERLSVDYGKSKLEFSIYPAPQVSTA
motub1/1-451	RIRKLTACSGLQGFLIFHSFGGGTSGSFTSLLMERLSLDYGKSKLEFAIYPAPQVSTA
hamtub1/1-451	RIRKLADQCTGLQGFLVFHSFGGGTSGSFTSLLMERLSVDYGKSKLEFSIYPAPQVSTA
rattub1/1-451	RIRKLADQCTGLQGFLVFHSFGGGTSGSFTSLLMERLSVDYGKSKLEFSIYPAPQVSTA
chicktub1/1-451	RIRKLADSCSGLQGFLIFHSFGGGTSGSFTSLLMERLSVEYGKSKLEFAIYPAPQASSA
frogtub1/1-451	RIRKLADQCTGLQGFLVFHSFGGGTSGSFTSLLMERLSVDYGKSKLEFAIYPAPQVSTA
Htub1/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
Htub2/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
ptub1/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
ptub2/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
motub1/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
hamtub1/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
rattub1/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
chicktub1/1-451	VVEPYNSVLTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA
frogtub1/1-451	VVEPYNSILTTHTTLEHSDCAFMDNEAIYDICRRNLDIERPTYTNLNLISQIVSSITA

Figure 12

Htub1/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLVITYAPIISA EKAYHEQLSVAEITSSCFEPNS
Htub2/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLATYAPVISA EKAYHEQLSVADITNACFEPAN
ptub1/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLATYAPVISA EKAYHEQLSVAEITNACFEPAN
ptub2/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLATYAPVISA EKAYHEQLSVAEITNACFEPAN
motub1/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLVITYAPIISA EKAYHEQLSVAEITSSCFEPNS
hamtub1/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLATYAPVISA EKAYHEQLTVAEITNACFEPAN
rattub1/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLATYAPVISA EKAYHEQLSVAEITNACFEPAN
chicktub1/1-451	SLRFDGALNVDLTEFQTNLVPFPRIHFPLVITYAPIISSDRAYHEQLSVAEITSSCFEPNN
frogtub1/1-451	SLRFDGALNVDLTEFQTNLVPYPRIHFPLATYAPVISA EKAYHEQLTVADITNACFEPAN
Htub1/1-451	QMVKCDPRHGKYM ACCMLYRGDVVPKDVNVAIAAIAIKTKRTIQFVDWCPTGFKVGINYP
Htub2/1-451	QMVKCDPHGKYM ACCLLYRGDVVPKDVNAIAIATIKTKRTIQFVDWCPTGFKVGINYP
ptub1/1-451	QMVKCDPRHGKYM ACCLLYRGDVVPKDVNAIAAIAIKTKRSIQFVDWCPTGFKVGINYP
ptub2/1-451	QMVKCDPRHGKYM ACCLLYRGDVVPKDVNAIAIATIKTKRTIQFVDWCPTGFKVGINYP
motub1/1-451	QMVKCDPRHGKYM ACCMLYRGDVVPKDVNVAIAAIAIKTKRTIQFVDWCPTGFKVGINYP
hamtub1/1-451	QMVKCDPRHGKYM ACCLLYRGDVVPKDVNAIAIATIKTKRTIQFVDWCPTGFKVGINYP
rattub1/1-451	QMVKCDPRHGKYM ACCLLYRGDVVPKDVNAIAIATIKTKRTIQFVDWCPTGFKVGINYP
chicktub1/1-451	QMVKCDPQQGKYM ACCMLYRGDVVPKDVNVAIAAIAIKTNRSLQFVDWCPTGFKVGINYP
frogtub1/1-451	QMVKCDPRHGKYM ACCLLYRGDVVPKDVNAIAIATIKTKRSIQFVDWCPTGFKVGINYP
Htub1/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
Htub2/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
ptub1/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
ptub2/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
motub1/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
hamtub1/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
rattub1/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
chicktub1/1-451	IPTPGGDLAQVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVSEGMEGEFSE
frogtub1/1-451	TVVPGGDLAKVQRAVCMLSNTTAIAEAWARLDHKFDLMYAKRA FVHWYVGEGMEEGEFSE
Htub1/1-451	AREDLAALEKDYEEVGTDSEFEE--NEGEEF
Htub2/1-451	AREDMAALEKDYEEVGVDSEGEGEEEGEY
ptub1/1-451	AREDMAALEKDYEEVGIDSYEDE--DEGEE-
ptub2/1-451	AREDMAALEKDYEEVGVDSEGEGEEEGEY
motub1/1-451	AREDLAALEKDYEEVGTDSEFEE--NEGEEF
hamtub1/1-451	AREDMAALEKDYEEVGADSAEGD--DEGEY
rattub1/1-451	AREDMAALEKDYEEVGVDSEGEGEEEGEY
chicktub1/1-451	AREDLAALEKDYDEVATDLFEDE-NEAGDS-
frogtub1/1-451	AREDMAALEKDYEEVGADSADAE--DEGEY

Figure 12 (continued)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
10 June 2004 (10.06.2004)

PCT

(10) International Publication Number  
**WO 2004/047615 A3**

(51) International Patent Classification<sup>7</sup>: **C07C 247/12**,  
229/36, G01N 33/533, 33/534, 33/567

(21) International Application Number:  
PCT/US2003/037393

(22) International Filing Date:  
21 November 2003 (21.11.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/428,050 21 November 2002 (21.11.2002) US

CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,  
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,  
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **WYETH**  
[US/US]; Five Giralda Farms, Madison, NJ 07940 (US).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **GREENBERGER**,  
Lee, M. [US/US]; \*.

(74) Agents: **FEHLNER, Paul, F.** et al.; Darby & Darby P.C.,  
Post Office Box 5257, New York, NY 10150-5257 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,

**Published:**


- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(88) Date of publication of the international search report:  
12 August 2004

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: **HEMIASTERLIN AFFINITY PROBES AND THEIR USES**

(57) Abstract: Photoaffinity probes are provided that are based hemiasterlin and derivative compounds thereof. Use of these probes to identify binding sites for these and other drugs, particularly anti-tubulin drugs, are also provided as are methods for identifying new drugs (*e.g.*, new anti-tubulin drugs) that bind to these binding sites.



**WO 2004/047615 A3**

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/37393

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07C 247/12, 229/36; G01N 33/533, 33/534, 33/567

US CL : 552/11; 562/441; 436/501, 503, 545, 546, 815

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 552/11; 562/441; 436/501, 503, 545, 546, 815

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,661,175 A (KASHMAN et al) 26 August 1997 (26.08.1997), Compounds 2 and 3.	1-29
A	WO 99/32509 A2 (THE UNIVERSITY OF BRITISH COLUMBIA) 01 July 1999 (01.07.1999), Compound I.	1-29
A	US 6,153,590 A (ANDERSEN et al) 28 November 2000 (28.11.2000), Compound I.	1-29
A	GAMBLE, W.R. et al. Cytotoxic and Tubulin-Interactive Hemiassterlins from Auletta sp. and Siphonochalina spp. Sponges. Bioorganic & Medicinal Chemistry. 1999, Vol. 7, pages 1611-1615, especially Compounds 1-4.	1-29

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

01 June 2004 (01.06.2004)

Date of mailing of the international search report

02 JUL 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Mary E. Ceperley

Telephone No. (571) 272-1600

**INTERNATIONAL SEARCH REPORT**

PCT/US03/37573

**Continuation of B. FIELDS SEARCHED Item 3:**

APS, CAS ONLINE

search terms: structure searches, hemiasterlin, anti-tubulin, tubulin binding, fluorography, photolabel, photoaffinity, photoreactive, benzophenone, azide, affinity

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**